



Exceptional Association of a Common Variable Immunodeficiency and Ankylosing Spondylitis: A Case Report

H. Bellamine ^{a*}, M. Moudatir ^a, K. Echchilali ^a, L. Barakat ^a,
M. Benzakour ^a and H. Elkabli ^a

^a *Internal Medicine Department, CHU Ibn Rochd, Casablanca, Morocco.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/117006>

Case Study

Received: 04/03/2024

Accepted: 08/05/2024

Published: 14/05/2024

ABSTRACT

We report a new observation of an unusual association of CVID with ankylosing spondylitis (AS). Osteoarticular involvement associated with CVID occurs in approximately 5 to 40% of cases, most commonly in the form of septic arthritis, or non-septic arthritis, which is part of the joint manifestations of autoimmune and rheumatic diseases associated with CVID. A 53-year-old patient with a history of recurrent bronchopulmonary infections complicated by bronchial dilatation since the age of 38, pelvic-axial syndrome and peripheral inflammatory arthralgia affecting the large and medium-sized joints and enthesitis (talalgia), who has presented since the age of 40 with a digestive disorder consisting of chronic liquid diarrhoea with 6 stools a day, developing in the context of a deterioration in her general condition, without any mention of medication. Regarding biotherapy, in particular anti-TNF alpha, this is the first-line biotherapy for APS. It has been shown to be effective in patients with CVID with joint involvement in the course of granulomatous disease, which would be beneficial for our patient.

*Corresponding author: E-mail: hindbellamine@gmail.com;

Cite as: Bellamine, H., Moudatir, M., Echchilali, K., Barakat, L., Benzakour, M., & Elkabli, H. (2024). Exceptional Association of a Common Variable Immunodeficiency and Ankylosing Spondylitis: A Case Report. *Asian Journal of Case Reports in Medicine and Health*, 7(1), 64–67. Retrieved from <https://journalajcrmh.com/index.php/AJCRMH/article/view/188>

Keywords: Ankylosing spondylitis; granulomatous diseases; immunoglobulins; autoimmune manifestations.

ABBREVIATION

AS: Ankylosing Spondylitis

1. INTRODUCTION

Common variable immunodeficiency (CVID) is a rare disease. It is the most common constitutional humoral deficiency of CVID in adults, characterised by polymorphic manifestations, in particular a decrease in the serum concentration of most circulating immunoglobulins (Ig), and by the occurrence of recurrent bacterial infections, autoimmune manifestations, malignant tumours or malabsorption.

Osteoarticular involvement associated with CVID occurs in approximately 5 to 40% of cases, most commonly in the form of septic arthritis or non-septic arthritis, which is part of the joint manifestations of autoimmune and rheumatic diseases associated with CVID. Bone involvement is less common.

The discovery of hypogammaglobulinemia in inflammatory rheumatism is a rare event.

We report a new observation of an exceptional association of CVID with ankylosing spondylitis (AS).

2. CASE PRESENTATION

A 53-year-old male patient with a history of recurrent bronchopulmonary infections since the age of 38, complicated by bronchial dilatation, pelvic-axial syndrome and peripheral inflammatory arthralgia of the large and medium-sized joints and enthesitis (talalgia), who presented with a digestive disorder since the age of 40, consisting of chronic liquid diarrhoea with 6 stools per day, developing in the context of a deterioration in his general condition, without any mention of medication. On clinical examination, the patient was in relatively good general condition, with bilateral snoring and achromatic patches on the anterior surfaces of both legs, consistent with vitiligo. Paraclinical investigations revealed an inflammatory syndrome with a CRP of 27mg/l, and the CBC showed a hyperleukocytosis of 14400/mm³ with a predominance of PNN (70%). Stool tests for bacteria and parasites were negative. HIV serology was negative. Protein electrophoresis showed a significant hypogammaglobulinemia of

0.6 g/l (Fig. 1), and low levels of IgG1 (<0.15), IgG2 (<0.02 g/l) and IgA (<0.007 g/l) by weight. Examination of lymphocyte subpopulations showed no qualitative or quantitative abnormalities. Rheumatoid factor, anti-CCP and ANA were negative. IgG and IgA anti-transglutaminase antibodies were negative, gastric fibroscopy and biopsy showed partial duodenal villous atrophy, chest CT showed bilateral bronchial dilatation, HLA b27 was negative, and CT of the sacroiliac joints showed bilateral stage III sacroiliitis (Fig. 2). A diagnosis of digestive CVID with systemic autoimmune manifestations such as vitiligo associated with ankylosing spondylitis (AS) was accepted.




Fig. 1. Serum protein electrophoresis: profound hypogammaglobulinemia



Fig. 2. Subchondral condensation of the joint edges: sacroiliitis III

The treatment regimen consisted of a monthly immunoglobulin infusion of 0.8 g/kg, antibiotic prophylaxis, and corticosteroid therapy of 10 mg/d with a slow taper of the corticosteroids,

maintaining a low dose of 5 mg/d as maintenance treatment in conjunction with the monthly intravenous infusion.

The therapeutic progress was notable, particularly in the osteoarticular and digestive areas.

3. DISCUSSION

Primary immune deficiencies (PIDs) are defined as functional and/or quantitative abnormalities in one or more components of the immune system, including T or B lymphocytes, natural killer cells, phagocytic cells, complement proteins, and others.

Ankylosing spondylitis (AS) is a chronic inflammatory rheumatic disease that primarily affects the axial skeleton and the sacroiliac joint, but can also involve peripheral joints and entheses. Following the discovery of the human leukocyte antigen B27 (HLA-B27), several hypotheses have been put forth to elucidate the precise etiology of AS. The tendency of HLA-B27 to form unusual structures leads to the recognition and activation of crucial components of the innate immune system. In recent years, SpA has been classified as a polygenic autoinflammatory disease, in which innate immune abnormalities may play a significant role [1,2].

These PIDs are manifested by repeated infections, which typically occur during childhood. However, other clinical manifestations are also possible. Some are highly specific and rare, while others are more frequent, particularly osteoarticular complications. These are caused by a wide variety of mechanisms, and are observed in different types of DIP, mainly humoral [3]. The prevalence of osteoarticular involvement in CVID is estimated to range from 5 to 40% [4].

Although bone manifestations are relatively uncommon, arthritis is a more prevalent condition. These are arthralgias, which may be genuine septic or aseptic. They may resemble rheumatoid arthritis but never ankylosing spondylitis [5].

The occurrence of hypogammaglobulinemia in the context of inflammatory rheumatism is a relatively uncommon phenomenon, which presents a diagnostic challenge that is frequently challenging to resolve. In practice, several hypotheses have been proposed to explain this phenomenon.

The condition may be the result of an inflammatory rheumatism that has been complicated by an immune deficiency secondary to background treatment or corticosteroid therapy. This is the most common occurrence. However, this was not the case in our observation, as the patient had not previously received any treatment that caused CVID, and the diagnosis of SPA was made at the same time as that of immune deficiency.

An alternative hypothesis is that the patient may be suffering from a genuine humoral DIP. However, this is a rare or even exceptional situation, given that these arthropathies are not associated with enthesopathy and do not, apart from exceptional cases, progress to an APS picture. Furthermore, there is no established link with HLA-B27 [6].

Alternatively, it may be a lymphoproliferative syndrome complicated by an immune deficiency, which is revealed by joint manifestations.

This presents a significant challenge in the treatment of ankylosing spondylitis. We were hesitant to prescribe anti-inflammatory drugs due to the patient's digestive involvement, in order to avoid exacerbating the condition.

Nevertheless, efficacious pharmaceutical agents have yet to be developed, and a significant proportion of patients encounter difficulties or become incompatible with existing pharmacological treatments [7].

Although corticosteroid therapy can induce moderate hypogammaglobulinemia, it was the only treatment the patient received that demonstrated a statistically significant improvement in joint damage at a rate of 5 mg/d. It would appear that the infectious consequences are low [8].

In contrast to immunosuppressive drugs such as methotrexate, which induce neutropenia and therefore indirectly facilitate bacterial infections, corticosteroids do not appear to have this effect.

With regard to biotherapy, in particular anti-TNF alpha, this is the recommended first-line biotherapy for APS. It has been demonstrated to be efficacious in patients presenting with CVID and joint involvement in the context of granulomatous diseases, a scenario that would be advantageous for our patient. Nevertheless, this form of treatment should be employed with caution, given the increased risk of infectious complications, the most significant of which is the reactivation of tuberculosis [9].

It is therefore advisable to exercise caution when considering the use of these therapies, provided that antibiotic prophylaxis has been initiated.

Intravenous immunoglobulin (IVIg) has been demonstrated to be efficacious in the treatment of aseptic arthritis in patients with common variable immunodeficiency (CVID), although there is no consensus on this matter. Moreover, there is a paucity of studies in the field of AS [10,11].

4. CONCLUSION

In conclusion, the occurrence of hypogammaglobulinemia associated with inflammatory rheumatism is a relatively uncommon scenario, often difficult to diagnose, and may occur secondary to immunosuppressive therapies. However, our observation represents a unique case of a true common variable immunodeficiency disorder (CVID) associated with ankylosing spondylitis (AS), a combination not previously reported in the medical literature.

The therapeutic management of such cases is complicated by the limited treatment options available due to the increased risk of infection in individuals predisposed to hypogammaglobulinemia.

CONSENT

As per international standards or university standards, patient(s) written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Vanaki N, Aslani S, Jamshidi A, Mahmoudi M. Role of innate immune system in the

- pathogenesis of ankylosing spondylitis. *Biomedicine & Pharmacotherapy*. 2018 Sep 1;105:130-43.
2. Díaz-Peña R, López-Vázquez A, López-Larrea C. Old and new HLA associations with ankylosing spondylitis. *Tissue antigens*. 2012 Sep;80(3):205-13.
3. *Revue du Rhumatisme*. Bone and joint disease associated with primary immune deficiency. Christelle Sordet. 2005; 72:1270–1282. DOI:10.1016/j.rhum.2004.07.015.
4. Hansel TT, Haeny MR, Thompson RA. Primary hypogammaglobulinemia and arthritis. *BMJ* 1987;295:174–5.
5. Katherine A. al Osteoarticular infectious complications in patients with primary immune deficiency. DOI: 10.1097/BOR.0b013e3282fd6e70
6. Andres E, Limbach FX, Kurtz JE, Kurtz-Illeg V, Schaefferbeke T, Kuntz JL, et al. Primary humoral immunodeficiency (late-onset common variable immunodeficiency) with antinuclear antibodies and selective immunoglobulin deficiency. *Am J Med*. 2001;111:489–91.
7. Fei Y. al The causal relationship between immune cells and ankylosing spondylitis: A bidirectional Mendelian randomization study. *Arthritis Res Ther*. 2024 Jan 16;26(1):24. DOI: 10.1186/s13075-024-03266-0.
8. Maxime Samson et al. Diagnostic strategy for patients with hypogammaglobulinemia in rheumatology. *Joint Bone Spine*. 2011;78:241–245.
9. Smith KJ et al. Common variable immunodeficiency treated with a recombinant human IgG, tumour necrosis factor-alpha receptor fusion protein. *Br J Rheumatol*. 2001;144:597–600.
10. Ben Mulhearn et al. Indication for igiv in rheumatic diseases; 2015, Oxford.
11. Webster A D, et al. Polyarthritis in adults with hypogammaglobulinaemia and its rapid response to immunoglobulin treatment. *Br Med J*; 1976.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/117006>