



Mandibular Sialoadenectomy in Dog with Sialocele: Case Report

**Larissa Barbosa Torres ^a, Eliza Maria Dias ^a
and Iago Martins Oliveira ^{b*}**

^a *Veterinarian in the City of Goiânia, Brazil.*

^b *School of Medical and Life Sciences, Pontifícia Universidade Católica de Goiás, Brazil.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/air/2024/v25i41108>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/119761>

Case Report

Received: 16/05/2024

Accepted: 17/07/2024

Published: 18/07/2024

ABSTRACT

The report aims to describe a case of left mandibular sialocele in a dog. A male, castrated, Shih Tzu dog, weighing 7 kilograms and five years old, was treated at the Veterinary Hospital of the School of Veterinary and Zootecnics at the Federal University of Goiás. A fluctuating non-painful subcutaneous swelling was observed in the left ventral cervical region during the clinical examination. Image tests revealed anechoic content with dilation of the salivary duct, heterogeneity and enlargement of the left mandibular gland and possibility of sialolithiasis. Besides, cytological analysis showed results that confirm the diagnose of left mandibular sialocele. Following the diagnosis, surgical treatment was made, performing sialoadenectomy. A skin incision was made ventral to the ear canal in the dorsal direction caudal to the angle of the mandible, and the patient was evaluated for days. Salivary sialocele is the most common sialoadenopathy in dogs and its diagnoses require several image methods and the surgical treatment is essential for healing and improving clinical signs.

*Corresponding author: E-mail: iago.vetufg@gmail.com;

Cite as: Torres, Larissa Barbosa, Eliza Maria Dias, and Iago Martins Oliveira. 2024. "Mandibular Sialoadenectomy in Dog With Sialocele: Case Report". *Advances in Research* 25 (4):311-15. <https://doi.org/10.9734/air/2024/v25i41108>.

Keywords: Canine; surgery; salivary gland; salivary duct.

1. INTRODUCTION

Salivary sialoceles or mucocele consists of an accumulation of mucinous secretion resulting from the extravasation of saliva into the interstitial space adjacent to the glands and salivary ducts [1]. It is characterized by an increase in volume, with a smooth, regular surface, well defined, unilateral or bilateral, painless and flaccid. Furthermore, it has a different anatomical location that depends on the affected gland [2,3].

The etiology of salivary mucoceles may be related to infections, trauma, obstructions or rupture of ducts, sialoliths, neoplasms, in addition to the idiopathic form [4]. Most of the time it is asymptomatic, although some animals may present dysphagia, abnormal tongue movements and inappetence [4,5].

Diagnosis is based on anamnesis, physical examination, fluid aspiration cytology, radiography and ultrasound. Treatment can be carried out conservatively through periodic drainage or glandular excision [3,5]. In view of the above, the objective of the present work is to report the use of the sialoadenectomy technique to treat mandibular gland sialoceles in a dog.

2. PRESENTATION OF CASE

A male, castrated, Shih Tzu dog, weighing 7 kilograms and five years old, was treated at the Veterinary Hospital of the School of Veterinary and Zootecnics at the Federal University of Goiás. The owner's main complaint was the appearance of a non-progressive increase in volume in the left cervical region.

During clinical examination, physiological parameters were unchanged, and a fluctuating non-painful subcutaneous swelling was observed in the left ventral cervical region. Given the clinical board, mandibular sialoceles was suspected. Complementary exams were requested like cervical radiography and ultrasound, blood count and hepatic and renal biochemical profile through measurement of serum alanine aminotransferase (ALT), alkaline phosphatase (AF), urea and creatinine.

The ultrasound examination showed anechoic content with dilation of the salivary duct, heterogeneity and enlargement of the left mandibular gland. The radiographic examination ruled out the possibility of sialolithiasis and the other analyzes were within the reference range for the species. Subsequently, a cytological



Fig. 1. Dissection of the mandibular salivary gland with presence of sialocel



Fig. 2. Salivary gland with mucocele associated with healthy gland tissue

analysis was carried out on the liquid collected from the affected region, which had a hypocellular, mucinous, yellowish and viscous appearance. Based on the information obtained, the diagnosis of left mandibular sialoceles was established and the patient was referred for surgical treatment.

A skin incision was made ventral to the ear canal in the dorsal direction caudal to the angle of the mandible. Careful dissection of the tissues was carried out and hemostasis was performed using electrocoagulation and in larger vessels, suturing in a crossed mattress pattern with number 3-0 polyglactone thread. The muscular fascia was incised, which promoted glandular exposure. Subsequently, dissection of the mandibular and sublingual glands was carried out up to the lingual branch of the trigeminal nerve (Fig. 1), ductal ligation with number 3-0 polyglactone suture and transection of the duct-gland complex (Fig. 2). Finally, the region was washed with 0.9% warmed saline solution. For synthesis, the subcutaneous tissue was reduced with number 3-0 polyglactone thread in an intradermal pattern and the dermorrhaphy was performed with number 3-0 nylon thread in a separate simple pattern. A passive drainage system with a Penrose drain was implanted in the ventral aspect of the surgical wound.

On the fourth postoperative day, the drain was removed and 10 days after the procedure, the skin stitches were removed. Reevaluation was made after 60 days, and the patient did not present a recurrence.

3. DISCUSSION

The patient in the present report presented a mucocele of the left mandibular gland, in line with the description [5-8] of a higher prevalence of cervical sialoceles in canine patients. It was not possible to determine the cause, mainly because the patient described here had no history of previous trauma and the cervical x-ray ruled out lithiasis, in contrast to the statement of a greater association of this disease with obstructions or ruptures secondary to sialoliths [4,9].

The sialodocentesis performed was compatible with material poor in cellularity, with a mucinous and cloudy appearance. However, cytological analysis normally demonstrates vacuolated macrophage infiltrate, coalescent and polymorphonuclear cells [10]. The ultrasound findings in this case are similar to those described by Kealy et al. [11] for sialoceles characterized by an area with anechoic content and absence of flocculation.

The treatment performed in this case was sialoadenectomy of the mandibular and lingual glands, which corroborates the indication for choosing this surgical modality [3,5,10]. The choice of sialoadenectomy in relation to other treatments, such as drainage, aspiration and drug treatments, is due to the surgical approach being definitive for sialoceles, preventing the condition from recurring. Furthermore, there is a reduction in the occurrence of complications such as secondary infections, worsening the patient's prognosis [5]. There were pre and intra-operative complications and tissue division was meticulously performed with special care to avoid accidental incision of the glandular capsule and leakage of contents at the surgical site. It is known that salivary gland excision is a simple, quick, effective procedure and, when there's no saliva contamination infections are rare [2,12]. The technique used was associated with a passive postoperative drainage system with removal after four days, as well as Penrose or negative pressure drains [5]. No suture dehiscence or seroma was observed, unlike some reported cases in which xerostomia, interference with swallowing and a large amount of seroma were observed [5,13,14].

4. CONCLUSION

The sialoadenectomy technique is effective in correcting mandibular gland sialoceles and allows effective removal of the duct-gland complex.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

CONSENT

It is not applicable.

ETHICAL APPROVAL

Animal Ethic committee approval has been collected and preserved by the author(s)

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Adams P, Halfacree J, Lamb CR. Zygomatic salivary mucocoele in a Lhasa apso following maxillary tooth extraction. *The Veterinary Record*. 2011;168(3):458-462.
2. Vallefucio R, Jardel N, Elmrini M, Stambouli F, Cordonnier N. Parotid salivary duct sialoceles associated with glandular duct stenosis in a cat - case report. *Journal of Feline Medicine and Surgery*. 2011;13(1):781-783.
3. Kazemi D, Doustar Y, Assadnassab G. Surgical treatment of a chronically recurring case of cervical mucocele in a German shepherd dog. *Case Reports in Veterinary Medicine*. 2012;1(1):1-4.
4. Gahir D, Clifford N, Yousefpour A, Avery C. A novel method of managing persistent parotid sialoceles. *British Journal of Oral and Maxillofacial Surgery*. 2011;49(1):491-492.
5. Fossum TW. Digestive system surgeries. In: Radlinsky MG. *Small animal surgery*: 4th ed. São Paulo: Roca; 2014;222-405.
6. Ritter MJ, Stanley BJ. Salivary glands. In: Tobias KM, Johnston SA, eds. *Veterinary Surgery, Small Animal*. 2nd ed. St Louis, MO: Elsevier/Saunders. 2018: 1653-1663.
7. Poirier VJ, Mayer-Stankeová S, Buncholz J, Vail DM, Kaser Hotz B. Efficacy of radiation therapy for the treatment of sialoceles in dogs. *J Vet Intern Med*. 2018;32:107-110.
8. Ritter MJ, von Pfeil D, Stanley B, Hauptman JG, Walshaw R. Mandibular and sublingual sialoceles in the dog: a retrospective evaluation of 41 cases, using the ventral approach for treatment. *N Z Vet J*. 2006;54:333-337.
9. Fernandes TR, Grandi F, Monteiro LN, Salgado BS, Rocha RM, Rocha NS. Ectopic ossification presenting as osteoid metaplasia in a salivary mucocele in a Shi Tzu dog. *BMC Veterinary Research*. 2012;8(13):2-5.
10. Andrade EC, Sepulveda RV, Galvão SR, Delcarlo RJ. Bilateral resection of salivary glands in the treatment of cervical sialoceles in a dog - case report. *CFMV Magazine – Federal Council of Veterinary Medicine*. 2011;17(54):44-48.
11. Kealy JK, Mcallister H, Graham JP. Skull and spine. In: Kealy JK, Mcallister H, Graham JP. *Radiography and ultrasound*

- of the dog and cat: 5th ed. Rio de Janeiro: Elsevier. 2012;447-453.
12. Cinti, F, Rossanese M, Buracco P, Pisani G, Vallefucio R, Massari F, Cantatore M. Complications between ventral and lateral approach for mandibular and sublingual sialoadenectomy in dogs with sialoceles. *Veterinary Surgery*. 2021;50(3): 579-587.
 13. Kaiser S, Thiel C, Kramer M, Pepler C. Complications and prognosis of cervical sialoceles in the dog using the lateral surgical approach. *Tierarztl Prax Ausg K Kleintiere Heimtiere*. 2016;44:323-333.
 14. Tsioli V, Papazoglou LG, Basdani E. Surgical management of recurrent cervical sialoceles in four dogs. *J Small Anim Pract*. 2013;54:331-333.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/119761>