



# **Microbiological Profile of Neonatal Septicemia by Automated Blood Culture System (BacT/ALERT) at a Tertiary Care Hospital in Jamnagar, Gujarat, India**

**J. Ghedia Bindiya<sup>a#</sup> and Manish R. Shah<sup>b\*≡</sup>**

<sup>a</sup> Department of Microbiology, Smt B. K. Shah Medical Institute & Research Centre, Sumandeep Vidyapeeth Deemed to be University, Piparia 391760, Vadodara, Gujarat, India.

<sup>b</sup> Department of Orthopaedics, Dhiraj Hospital, Smt. B. K. Shah Medical Institute & Research Centre, Sumandeep Vidyapeeth Deemed to be University, Piparia, Vadodara, 391760, Gujarat, India.

## **Authors' contributions**

*This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/JPRI/2021/v33i59B34346

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/79795>

**Original Research Article**

**Received 10 October 2021**  
**Accepted 14 December 2021**  
**Published 17 December 2021**

## **ABSTRACT**

**Background:** This study was carried out to do rapid diagnosis of neonatal septicemia to prevent septicemia related complications and neonatal morbidity and mortality.

**Materials and Methods:** This study was carried out in a tertiary care hospital from a period of March 2014 to August 2015. This is a non-randomized, prospective study in which one hundred and nine cases of suspected neonatal septicemia on the basis of antenatal high risk factors and signs and symptoms of sepsis were studied. Blood samples were collected from neonates in whom septicemia was suspected usually before antibacterial agents were given and both positive and negative samples were detected by automated system BacT/ALERT.

**Results:** Out of 109 samples tested, 69 samples were positive and 40 samples were negative. The incidence of neonatal septicemia due to Gram positive organisms (62.85%) was found to be high. Among them *Coagulase negative staphylococci* (CONS) (27) (38.57%) was most common organism followed by *Staphylococcus aureus* (11) (15.71%), *Enterococcus spp.* (7.14%) and *Streptococcus spp.* (1.43%). Rate of Gram negative organisms was quite low (32.85%). Out of

<sup>#</sup> Microbiology Tutor;

<sup>≡</sup> Associate Professor;

\*Corresponding author: E-mail: manishshah2001in@yahoo.co.in;

them *Acinetobacter spp.* (17.14%) was most common followed by *Klebsiella spp.* (7.14%), *E.coli* (5.71%) and *Pseudomonas spp.* (2.86%).

**Conclusion:** Early detection of infection in neonates and proper antibiotics for the treatment leads to reduced morbidity and mortality.

**Keywords:** Neonatal septicemia; BacT/ALERT; gram positive organisms; gram negative organisms.

## 1. INTRODUCTION

Neonatal septicemia is a clinical syndrome that occurs due to acute invasion of micro-organisms and their active multiplication in the blood stream that has been documented by positive blood culture during the first month of life [1].

It is estimated that 20% of all neonates develop septicemia [2,3] and it is responsible for 30-50% of total deaths during the first month of life in developing countries [2,4]. Neonatal septicemia is the commonest cause of neonatal mortality in the developing countries like India [5], so prompt detection of micro-organisms through blood culture and appropriate antimicrobial therapy with the help of other laboratory investigations and supportive care are the key determinants for positive outcome in this serious pediatric emergency.

The fulminant nature of this illness makes it necessary to review the relative frequency of various bacteria responsible for this serious condition so that appropriate antibiotic treatment and patient care can be instituted as early as possible [6].

The reported incidence of neonatal septicemia in developed countries varies, so it is difficult to assess the true incidence in our country because of unavailability of community data and difficulty in getting microbiological studies from all hospitals in our country (as it may be only on clinical grounds).

Incidence of neonatal septicemia in hospital born infants in India is in range of 11.0-24.5 per 1000 live births [7].

The purpose of this study was to find out the etiological agents present in neonatal septicemia patients as early as possible by the automated blood culture system named BacT/ALERT(Biomérieux) .

## 2. MATERIALS AND METHODS

This study was carried out in a tertiary care hospital (Guru Govindsingh Hospital, Jamnagar,

Gujarat, India; 22.4707°N, 70.0577°E), from a period of March 2014 to August 2015. This is a non-randomized, prospective study in which one hundred and nine cases of suspected neonatal septicemia on the basis of antenatal high risk factors and signs and symptoms of sepsis including maternal fever, prolonged rupture of membranes, foul smelling lochia, temperature instability, feeding difficulty, respiratory distress, jaundice, convulsions, autonomic disturbances during this period were studied. Blood samples were collected from neonates in whom septicemia was suspected usually before antibacterial agents were given.

### 2.1 Inclusion Criteria

- i. Neonates (age – 0 to 28 days).
- ii. Neonates having signs and symptoms of septicemia such as maternal fever, feeding difficulty, convulsions and respiratory distress syndrome.
- iii. Premature infants.
- iv. Only blood samples of neonates and not any other fluid.

### 2.2 Exclusion Criteria

- i. Infants and children above the age of 28 days and adults and elderly patients.
- ii. Any other body fluid except blood.

Clinical history, birth history including treatment given was noted in all cases. All the details were taken within a proforma.

Blood samples were collected with disposable syringe and needle under strict asepsis. Usually samples were collected from peripheral vein. Skin was painted with 70% alcohol and allowed to dry. Then approximately 2 ml of blood was collected and immediately transferred to BacT/ALERT PF PLUS aerobic blood culture bottles. The bottles were incubated in BacT/ALERT system and patient's clinical data was introduced in the computer.

### 2.3 Principle of the Test

The BacT/ALERT microbial detection system utilizes a colorimetric sensor and reflected light to

monitor presence and production of carbon dioxide (CO<sub>2</sub>) dissolved in culture medium. If microorganisms are present in test sample, carbon dioxide is produced as the microorganisms metabolize the substrates in culture medium. When growth of the microorganisms produces CO<sub>2</sub>, the color of gas permeable sensor installed in the bottom of each culture bottle changes from blue-green to yellow. The lighter color results in an increase of reflectance units monitored by the system. Bottle reflectance is monitored and recorded by the instrument every 10 minutes.

Bottles that do not become positive remain in the system for 5 days. After 5 days, the computer prints out a list of negative bottles and illuminates the light adjacent to each well containing a negative bottle. These bottles are removed from the system and discarded and declared negative.

The positive bottles are unloaded and subcultures are made on macConkey's medium and blood agar plates. Organisms are identified from their colony characteristics, appearances in smear, biochemical reactions, appearance on mannitol salt agar and bile esculin agar.

### 3. RESULT

A total 109 samples were tested out of which 69(63.30%) were positive and 40(36.70%) were negative for neonatal septicemia. Out of 109 total cases, 69(63.3 %) were suspected of neonatal septicemia between 0-3 days of age and 40

(36.7%) were suspected between 4-28 days of age [Table 1].

Out of 109 cases of septicemia, 66 (60.55%) cases were suspected in male neonates, while 43(39.45%) were being suspected in female neonates [Table 2].

Out of 69 positive cases, 44 (63.8%) cases were noted positive between 0-3 days of age while 25 (36.2%) cases were noted positive between 4-28 days of age [Table 3].

45 (65.22%) were male neonates while 24 (34.78%) were female neonates [Table 4].

Out of 69 positive cases, in one positive case two organisms were isolated. So total organisms isolated were 70. Out of 70 organisms isolated, Gram positive cocci are 44 (62.85%), Gram negative bacilli are 23 (32.85%) and *Candida non-albicans* are 3 (4.30%) [Table 5].

Amongst gram positive bacteria, 27 (38.57%) of coagulase negative *staphylococci* (CONS), 11 (15.71%) of *Staphylococcus aureus*, 5(7.14%) of *Enterococcus spp.* and 1(1.43%) of *Streptococcal spp.* isolated from blood cultures. Amongst gram negative bacteria, 12 (17.14%) of *Acinetobacter spp.*, 5(7.14%) of *Klebsiella spp.*, 4(5.71%) of *E.coli*, 2(2.86%) of *Pseudomonas spp.* and amongst *Candida*, 3(4.30%) of *Candida nonalbicans* isolated from blood culture samples[ Table 6].

**Table 1. Distribution of cases according to age of onset of septicemia**

Age	Total no. of Cases	Percentage
0 - 3 days of age	69	63.3%
4-28 days of age	40	36.7%
Total	109	100%

**Table 2. Distribution of cases according to sex**

Sex	Total no. of Cases	Percentage
Male	66	60.55%
Female	43	39.45%
Total	109	100%

**Table 3. Distribution of positive cases according to age of onset of septicemia**

Age	Positive Cases	Percentage
0-3 days of age	44	63.8%
4-28 days of age	25	36.2%
Total	69	100%

**Table 4. Distribution of positive cases according to sex**

Sex	Positive Cases	Percentage
Male	45	65.22%
Female	24	34.78%
Total	69	100%

**Table 5. Organism wise distribution of positive cases of neonatal septicemia**

Organism	No. of cases	Percentage(%)
Gram positive cocci	44	62.85%
Gram negative bacilli	23	32.85%
<i>Candida nonalbicans</i>	3	4.30%
Total	70	100%

**Table 6. Distribution of positive cases according to individual organism**

Organism	No. of cases	Percentage
<b>Gram Positive Organism</b>		
1. <i>Coagulase negative staphylococci</i>	27	38.57%
2. <i>Staphylococcus aureus</i>	11	15.71%
3. <i>Enterococcus spp.</i>	5	7.14%
4. <i>Streptococcus spp.</i>	1	1.43%
<b>Total</b>	<b>44</b>	<b>62.85%</b>
<b>Gram Negative Organism</b>		
1. <i>acinetobactor spp.</i>	12	17.14%
2. <i>klebsiella spp.</i>	5	7.14%
3. <i>E.coli</i>	4	5.71%
4. <i>pseudomonas spp.</i>	2	2.86%
<b>Total</b>	<b>23</b>	<b>32.85%</b>
<b>Candida</b>		
1. <i>Candida nonalbicans</i>	3	4.30%
<b>Total</b>	<b>70</b>	<b>100%</b>

#### 4. DISCUSSION

Varieties of microorganisms like Gram positive and Gram negative bacteria and fungi like candida are responsible for neonatal septicemia. The most difficult tasks faced by neonatologist is to differentiate between septicemic and non-septicemic cases, this situation is complicated by conditions like birth asphyxia, hypoglycemia, hypothermia, intracranial hemorrhage. For definitive diagnosis of septicemia, causative microbes are required to be isolated from blood culture which takes minimum time of 48 hours. It also provides correct guideline to institute rational antibiotic therapy.

Present study was undertaken with the idea to obtain prevalent etiological agents and their rapid detection by BacT/ALERT system. So we can give appropriate antibiotics and reduce morbidity and mortality.

Neonatal septicemia must be treated urgently and specifically to reduce the mortality.

Suspected infection based on clinical criteria need to be supported by microbial investigations to find causative organisms and their antibiotic sensitivity pattern.

In present study rate of gram positive organisms was 62.85% which is comparable with the study of Kavitha Prabhu et al. (64.19%) [8] and Abhishek Mehta et al. (67.8%) [9] and Gram negative organisms were 32.85% which is comparable with the study of Kavitha Prabhu et al. (34.56%) [8] and Abhishek Mehta et al (28.7%) [9] and *Candida spp.* were 4.30% comparable with the Abhishek Mehta et al (2.87%) [9] and R. Nandana et al. (6.25%) [10].

A total number of 109 neonates with clinically suspected neonatal septicemia were studied. 69 (63.3%) cases out of them were blood culture positive which is comparable with the study of Izeta softic et al. [11]. Maximum number of neonatal septicemia was between 0-3 days of age (63.8%) which is similar to study done by

Vrishali et al. [12]. Neonatal septicemia occurs more in male patients (65.22%) than female patients (34.78%) which is similar to study done by Navdeep Saha et al. [13]. Early sepsis was more commonly associated with *coagulase negative staphylococci*, *acinetobacter species* and *staphylococcus aureus* whereas late sepsis was more commonly associated with *coagulase negative staphylococci*. In present study all samples show pure growth of organism. The incidence of neonatal septicemia due to gram positive organisms (62.85%) was found to be high. Similar findings of high positivity by gram positive organisms were seen in studies of Van den hoogen and Galhotra S [14,15] whereas some researchers (Muley et al 2015, Jyothi et al 2013) reported Gram negative organisms as predominant pathogens [16,17]. Among them *Coagulase negative staphylococci* 27 (38.57%) was most common organism followed by *Staphylococcus aureus* 11 (15.71%), *Enterococcus spp.* 5(7.14%) and *Streptococcus spp.* 1(1.43%) which is similar to study done by Bipin gupta et al. [18], Ballot DE et al. [19]. In studies done by other researchers [20,21] also CONS were the most common organisms isolated in neonatal septicemia cases. The increasing prevalence of (CONS) infections is due to their increasing antibiotic resistance as reported by some researchers [22]. Rate of gram negative organisms was quite low (32.85%). Out of them *Acinetobacter spp.* (17.14%) was most common followed by *Klebsiella spp.* (7.14%), *E.coli* (5.71%) and *Pseudomonas spp.* (2.86%). In other studies, other gram negative bacteria have been found to be most common gram negative bacteria e.g. *Klebsiella spp.* [23,24].

## 5. CONCLUSION

In summary, neonatal septicemia is more common during first 3 days of life and it is more common in male neonates. The Gram positive organisms are the frequent cause of neonatal septicemia. Early detection of neonatal sepsis along with broad spectrum antibiotics lead to reduced mortality rate. Better management of sepsis related complications have also reduced the mortality.

## DISCLAIMER

The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not

intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

## CONSENT

As per international standard, parental written consent has been collected and preserved by the author(s).

## ETHICAL APPROVAL

This study has been approved by ethical committee. After clearance of Institution Review Board (IRB) the study was conducted.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Martin RJ, Fanaraff AA, et al. Neonatal septicemia, Behrman's neonatal-perinatal medicine 3<sup>rd</sup> edition. 1983;650-656.
2. Jain Nk, Jain VM, Maheshwari S. Clinical profile of Neonatal Sepsis. Kathmandu University Medical Journal. 2003;1(2):117-120.
3. Gotoff SP. Neonatal Sepsis and meningitis: in: Nelson textbook of Paediatrics. (15<sup>th</sup> edition). Eds Behrman RE, Kleigman RM, Arbin AM.
4. Agrawal R, Sarkar N, Deorary AK and Paul VK. Sepsis in newborn. Ind. J. Paediatric Dec 2001; 68:1143-1147.
5. Singh M, Deorari AK, Paul VK, et al. Perinatal mortality in a hospital. Indian J. Med. 1991;94(B): 1-5.
6. Bhakoo ON, Aggarwal KC, Mohini C, Mahajan et al. Indi. Paedii. 1968;5:518-523.
7. Vinod K, Paul et al. Neonatal sepsis, Medical emergencies in children, 2<sup>nd</sup> edition. 93:115-118.
8. Kavitha Prabhu, Sevitha Bhat, and Sunil Rao. Bacteriologic Profile and Antibiogram of Blood Culture Isolates in a Pediatric Care Unit. J Lab Physicians. 2010;2(2):85-88.
9. Abhishek M Mehta, Navinchandra M. Kaore, Tukaram K. Prabhu. Microbial Profile of Neonatal septicemia in a tertiary

- care hospital of Bhopal November 2014. International Journal of Biomedical and Advance Research. 2014;5(10):499. DOI:10.7439/ijbar.v5i10.919
10. Nandana Reddy Jonnala R, Zion Eluzai, satyanarayana Rao N. Clinical and laboratory profile of neonatal sepsis.
  11. Izeta Softić, Husref Tahirović, Vincenzo Di Ciommo, Cinzia Auriti. Bacterial sepsis in neonates: Single centre study in a Neonatal intensive care unit in Bosnia and Herzegovina. Acta Med Acad 2017;46(1):7-15. DOI: 10.5644/ama2006-124.181
  12. Vrishali Avinash Muley, Dnyaneshwari Purushottam Ghadage, Arvind Vamanrao Bhore. Bacteriological profile of neonatal septicemia in a tertiary care hospital from Western India. Journal of Global Infectious Diseases. 2015;7(2):75-77.
  13. Navodeep Saha, Mallika Sengupta, Soma Sarkar and Manideepa Sengupta. Clinical and Microbiological Profile of Neonatal Septicemia in A Tertiary Care Hospital in Kolkata. J Pure Appl Microbiol. 2020;14(2):1537-1543. Article Number: 6162.
  14. Van den Hoogen A, Gerards LJ, Verboon Maciolek MA, Fleer A, Krediet TG. Long-term trends in the epidemiology of neonatal sepsis and antibiotic susceptibility of causative agents. Neonatology 2010;97:22-8.
  15. Galhotra S, Gupta V, Bains HS, Chhina D. Clinico-bacteriological profile of neonatal septicemia in tertiary care hospital. J Mahatma Gandhi Inst Med Sci 2015;20:148-52.
  16. Muley VA, Ghadage DP, Bhore AV. Bacteriological profile of neonatal septicemia in a tertiary care hospital from Western India. J Global Infect Dis 2015;7:75-7.
  17. Jyothi P, Metri C, Basavaraj, Peerapur V. Basavaraj, Bacteriological profile of neonatal septicemia and antibiotic susceptibility pattern of the isolates, Journal of Natural Science Biology and Medicine. 2013;4:306-309.
  18. Bipin Gupta, Sneha Mohan, Anjali Agarwal, Renu Dutta. Bacteriological Profile and Antibigram of Neonatal Septicemia in a Tertiary Care Hospital. 2018;7(8). ISSN: 2319-7706
  19. Ballot DE, Nana T, Sriruttan C, Cooper PA. Bacterial bloodstream infections in neonates in a developing country. ISRN Pediatr 2012;2012:508-512.
  20. Huaifu Dong, Huiping Cao and Haiyan Zheng. Pathogenic bacteria distributions and drug resistance analysis in 96 cases of neonatal sepsis.
  21. Shokry M, Bassyouni MI, Abu-El-Moon S, Maoz M, Tamer S. Evaluation of 16s rDNA amplification by PCR and some immunological mediators assessment compared with blood culture in diagnosis of neonatal sepsis. El-Minia Medical Bulletin. 2007;18:1-17.
  22. Ponce de Leon S, Wenzel RP. Hospital-acquired bloodstream infections with Staphylococcus epidermidis. Review of 100 cases. Am J Med. 1984;77:639-644.
  23. Narayan Gyawali, RK Sanjana. Bacteriological Profile and Antibigram of Neonatal Septicemia November 2012. The Indian Journal of Pediatrics. 2012; 80(5).
  24. K.V.Leela, R.Narayana Babu, Sugunya, Venkata Madhu Prasad and Ramisetty Raaga Deepa. Study of Bacterial Profile in Neonatal Sepsis and their Antibiotic Sensitivity Pattern in a Tertiary Care Hospital. 2016;5(6):511-521. ISSN: 2319-7706

© 2021 Bindiya and Shah; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/79795>