



Multiple Cocoon Crops Module and its Impact on Stabilization of Autumn Crop under Subtropical Condition of North West India

Sardar Singh ^{a*} and S. Murali ^b

^a Central Sericultural Research and Training Institute, Pampore-192121, J&K, India.

^b P2 BSF, NSSO, Central Silk Board, Horsley Hills, Kotevuru-517325, AP, India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2022/v12i121580

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/95161>

Original Research Article

Received: 20/10/2022

Accepted: 28/12/2022

Published: 29/12/2022

ABSTRACT

The present study was conducted mainly to explore the possibility of multiple cocoon crops in North West India and expected that introduction of the additional crops in between spring and the autumn will force the pruning of mulberry trees before autumn crop hence, it will help in productivity improvement during the autumn crop. The field trial was conducted with 9 selected rearers sponsored by Department of Sericulture, J&K & Himachal Pradesh and laboratory rearing was conducted at RSRS, Jammu. Three crop rearing schedule was followed with an innovative approach for date of brushing, leaf harvesting, silkworm rearing & mounting of larvae. The outcome of the study leads to understanding that change in brushing dates, rearing with shortlisted silkworm hybrids and use of improved rearing & mounting techniques can improve the cocoon productivity/100 DFLs in one hand and introduction of Summer crop in between spring & autumn crop can also improve the cocoon yield during the autumn crop. Farmer's level study revealed that

*Corresponding author: E-mail: dr.mmml@rediffmail.com;

an average yield recorded during spring crop of about 70.67 Kg & 60.33 Kg/100 DFLs in Jammu & HP whereas lab rearing at RSRS, Jammu recorded about 83.00 Kg/100 DFLs. During summer crop about 44.40 Kg & 35.35 Kg/100 DFLs in Jammu & HP and lab rearing at RSRS, Jammu recorded about 51.87 Kg/100 DFLs respectively. During autumn crop an average yield of about 40.80 Kg & 43.67 Kg/100 DFLs was recorded at Jammu & HP and lab rearing at RSRS, Jammu recorded about 44.75 Kg/100 DFLs. The impact of the introduction by summer crop revealed that *per cent* yield gain over traditional field rearing from experimental field trials about 67.14 (%) and 44.70 (%) yield gain records in the autumn season at Jammu division and Himachal Pradesh respectively.

Keywords: Multiple cocoon crops; Spring; Summer; Autumn; Jammu; Himachal Pradesh; Cocoon yield.

1. INTRODUCTION

One crop success story and the subsidiary nature of silk cocoon crops is a major limiting factor for sericulture development in North Western India. Though efforts were made except spring crop no other silk cocoon crop could be stabilized at the farmer level. Isolated efforts were made at the farm and farmer level but so far have not yielded desired results. Sericulture activities in the North-Western region of India are depending upon two crops- spring and autumn only, of which autumn crop is yet to be stabilized at farmer's level as a result about 75-80 (%) of the cocoon are being produced from spring crops only.

"Jammu & Kashmir, Himachal Pradesh, Uttarakhand, Punjab and Haryana contributes to the North Western sericulture zone of the country and are known for exclusively bivoltine cocoon production, no doubt it has good climatic condition during the spring rearing season for production of high quality bivoltine cocoons and there are very few pockets in the country which may be considered as a natural home for bivoltine silk. Hence over decades the entire zones depends on this crop for its annual raw silk output. There are only two silk cocoon crops in vogue in this zone *i.e.* spring and autumn of which spring crop alone constitutes about 80 *per cent* (except Uttarakhand) of the annual cocoon production" [1].

The important debated issue for the development of sericulture in North-West India is - can the number of silk cocoon crops will be increased, why the autumn crop is not stabilized at the farmer level and whether is there a scope for a third crop introduction between spring and autumn. It is evident from the recent cocoon production trends in the major cocoon producing states of North-West India [2]. During the year 2016-17 Jammu and Kashmir is the largest

cocoon producing state of North-West India with a production of about 973.30 MT of a bivoltine cocoon of which only 25.95 MT (2.66%) were produced during autumn crop [3]. Similarly in Himachal Pradesh the second largest cocoon producing state in the region, cocoon production during 2016-17 produced 236.55 MT of which only 35.40 (18%) were produced during the autumn crop [4]. Condition is slightly better in Uttarakhand which produced 231.37 MT of the cocoon of which 98.13 MT (42.41%) were produced during autumn crop. Yield /100 dfls during spring crop in J & K during 2016-17 was 39.41 Kg while during the autumn it was only 26.95 Kg while in Himachal Pradesh it was 41.01 Kg during spring and 26 Kg during the autumn [5]. The aim of the study was mainly to explore the possibility of multiple cocoon crops in North West India and expected that introduction of a crop between spring and the autumn will force the pruning of mulberry trees before the autumn crop hence, it will help in stabilization of autumn crop. Based on the study, the brushing dates and pruning schedule on three cocoon crop was worked out and an assessment was made for impact on the autumn crop through introduction of summer crops in between the spring & autumn crop.

2. MATERIALS AND METHODS

A preliminary work during 2018 was conducted at institute level investigation during 2019 to select suitable brushing dates for spring, summer and autumn crop and suitable hybrid combination was shortlisted for all three crops using the Mano index. Field trial were conducted by adopting three crops schedule with shortlisted hybrids with the farmers sponsored by the Department of Sericulture (DOS) in two states under given location covering 5 potential sericulture districts of North West India. Spring, Summer and autumn crop was conducted with shortlisted hybrid *i.e.* FC₁×FC₂ for spring crop & SH₆×NB₄D₂

for summer & autumn crop respectively with new brushing date at Jammu and Himachal region with selected silkworm rearers at each location with actual rearing capacity of the farmers. Chawki reared worms were distributed and leaves were used from the same trees for the three different seasons by adopting pruning from time to time (Table 1).

2.1 Methodology

Farmer's level (DOS sponsored farmers in potential sericulture districts) rearing was conducted with proven technology and innovative approach, based on the outcome of experimental rearing conducted at RSRS, Miran Sahib, Jammu during previous years (2018 & 2019). "Spring, summer and autumn field rearing was conducted during 2020 among DOS sponsored farmers (9 Nos) with shortlisted hybrid *i.e.* FC₁×FC₂ for spring crop & SH₆×NB₄D₂ for summer & autumn crop respectively (based on actual rearing capacity of the farmer) with overall technical and material support through RSRS, Jammu/State Sericulture Departments (DOSs) and Research Extension Centers (RECs) of CSB at the identified location. DOS, J&K and DOS, HP sponsored three and six farmers respectively, thus total 9 farmers with the help of the State Department of Sericulture/ Research Extension Centers of CSB in three different locations of representative sericulture districts in Jammu province of Jammu & Kashmir and Himachal Pradesh where finally identified. Accordingly farmer level rearing during spring, summer and autumn crop in J&K and HP was conducted with five potential sericulture districts - Rajouri, Udhampur & Kathua in J&K and district Bilaspur and Kangra in Himachal Pradesh comprising 03 farmers from each location having rearing capacity up to 50 dfls/farmer & Control rearing (50 DFLs) for the shortlisted hybrid was conducted at RSRS, Miran Sahib, Jammu to assess research level yield and potential yield at rearer's level. The introduction of summer crop in between spring and autumn was studied and its impact on the autumn crop was analyzed" [6]. The hybrid seed was procured from SSPC, Prem Nagar, Dehradun and were incubated at RSRS, Jammu and REC, Ghumarwin (H.P) worms were distributed after second moult for each selected rearers.

3. RESULTS AND DISCUSSION

The results of the field trial investigation at Jammu (J&K) and Himachal Pradesh with DOS

sponsored rearers during spring, summer and autumn (2020) seasons were presented here:

From the present study the results revealed that during spring rearing brushing was done on 01.03.2020 and the chawki reared worms were distributed to each location on 10.03.2020 having 50 DFLs per rearer. In the Jammu region of J&K cocoon yield is varied from 58 to 78 Kg /100 dfls. The actual yield recorded was about 76, 78 and 58 Kg/100 DFLs at Lamberi (Rajouri Dist.), Majalta (Udhampur Dist.) and Dharamakot (Kathua Dist.) regions respectively at Jammu division and the average yield was recorded as 70.67 Kg/100 DFLs whereas at Himachal Pradesh cocoon yield was in recorded from 36 to 74 Kg/100 dfls having an actual yield of 71, 36 and 74 Kg/100 DFLs at Soi, Panol and Naswal region of Bilaspur Dist., Himachal Pradesh and average yield was about 60.33 Kg/100 DFLs. The control batch reared at RSRS, Jammu under laboratory conditions achieved yield was about 83 Kg/100 DFLs (Table 2).

The result of the present study is in accordance with the results observed at North Kashmir during the summer season (2006) [7]. "Further, at the time of implementation of the cluster development program in Kashmir valley, summer crops were also undertaken in Tral and Bandipora clusters. The results obtained from cluster silkworm rearers were better [8] than the results obtained in North Kashmir" [7]. "A comparative study with 33 cluster and 30 non cluster silkworm rearers was conducted in the Tral area during summer season (2015) and it was revealed that the cluster farmers who have adequate infrastructure have harvested with an average cocoon yield of 33.72 Kg/100 DFLs whereas non cluster farmers have harvested 7.95 Kg/100 DFLs in the same area" [8] "The reasons behind the better results of cluster farmers may be the infrastructure developed under the programme and helped them to adopt the technologies as per the recommended package of practices" [8].

Similar results were in agreement with Singh and Singh [9] observed that successful trial of summer crop was conducted in Himachal Pradesh where 2000 dfls of PM×CSR2 was reared at farmer's level and an average yield of about 45 Kg/100 dfls was obtained during July 2010, where organized mulberry plantation was developed in 2006 were utilized for silkworm rearing.

Table 1. Showing pruning and brushing schedule adopted during different seasons of the study

Season	Pruning schedule	Brushing schedule	
		Jammu Condition (J&K)	H.P
Spring	Shoot harvesting for late age rearing during spring crop leaving behind a shoot bearing leaf on each branch.	Early first week of March (1 st of March)	Early 2 nd week of March (5 th -10 th March)
Summer	Shoot let harvesting for late age rearing during summer crop (June - July) adopting recommended method of monsoon pruning will forced the adoption of pruning practice in between spring and autumn crop which is absent at present and a pre-requisite for the success of autumn crop.	3 rd week of June (Between 15-30 th of June)	2 nd week of June (10-15 th of June)
Autumn	Apical harvesting of primary branches (above 3 feet crown on the main branch) during the autumn crop.	10-15 th of September	1 st week of September

Similarly a diagnostic study was conducted at Regional Sericultural Research Station Jammu in 2018 to explore the possible introduction of a third crop in between spring and autumn with a selection of suitable hybrid and brushing date for summer rearing under the Sub-tropical condition of Jammu. The hybrid selection was made after evaluating the pool of silkworm hybrid developed at different institutes. Brushing was ascertained after brushing the silkworm eggs at different dates and hybrids were evaluated according to the evaluation index for desired economic parameters for both quantity and quality developed by Mano et al. [10]. "Hybrids showing an index value above 50 were shortlisted. The first brushing was conducted on 10th of June 2018 with eight hybrids and the shortlisted hybrids were FC₁×FC₂, D×O₂, D×O₃ and SH₆×NB₄D₂ whereas Nine hybrids were reared during the second brushing of the summer season (2018), brushed on 25.06.2018. The hybrids showing an index value above 50 were shortlisted as PM×FC₂, PM×CSR2, SH₆×NB₄D₂, SK6×SK7 and D×O₃. Based on the study promising silkworm hybrid and suitable date of brushing for introduction of additional summer crop under sub-tropical condition has been workout" by Singh and Murali (2019).

"A new date of brushing has been assessed which suggest that the Spring crop in the region needs to be preponed by a week over the traditional date(s), Summer crop can be conducted in the month of June between the 2nd to 4th week and the autumn crop to be postponed by a week over the traditional date of brushing

especially in two major sericulture states *i.e.* Jammu region (Sub tropical) of J & K and Himachal Pradesh respectively" [1].

3.1 Impact on Autumn Crop by Introduction of Summer Crop

Measured in terms of *per cent* gain over existing practices after diffusions of knowledge & innovation and introduction of summer crop in between spring & autumn crop.

During summer crop the yield recorded about 44.40 Kg & 35.35 Kg/100 DFLs in Jammu & HP and lab rearing at RSRS, Jammu recorded about 51.87 Kg/100 DFLs and the average yield recorded by rearers at Jammu division of J&K during the autumn season was at about 40.80 Kg and HP was recorded about 43.67 Kg/100 dfls respectively. The *per cent* yield gain over traditional field rearing from experimental field trial revealed that 67.14 (%) and 44.70 (%) gain records in the autumn season at Jammu division and Himachal Pradesh in the same area respectively (Table 3).

"The summer crop is completed in the month of July last and fresh leaf is available for the next autumn crop after harvesting the leaf during summer rearing, it was a sort of forced pruning which otherwise is not done before the autumn crop. This had helps in the improvement of autumn silkworm crop performance and those farmers who are not opting for mulberry pruning after spring crop should take summer crop and sericulture becomes more remunerative" [6].

Table 2. Average data recorded during spring field trial for the hybrid, FC₁×FC₂ through shelf rearing (2020) at different places of Jammu & Himachal Pradesh and control rearing at RSRS, Jammu

Name of Location	Date of incubation	Date of brushing	H (%)	Date of distribution	Wt. of 10 Mature Larvae (g.)	5th Instar Larval Period (D:H)	Total Larval Period (D:H)	Actual Yield (Kg)	Av. Yield / 100 DFLs	SCW (g)	SSW (g)	SR (%)
Jammu Division (J&K)												
Lamberi, (Rajouri)	21.02.2020	01.03.2020	94.33	10.03.2020	55.00	6.00	29.00	38.00	76.00	1.90	0.40	21.04
Majalta, (Udhampur)	21.02.2020	01.03.2020	94.33	10.03.2020	56.00	6.19	31.00	39.00	78.00	1.763	0.393	22.29
Dharamkot, (Kathua)	21.02.2020	01.03.2020	94.33	10.03.2020	58.00	8.02	29.12	29.00	58.00	1.70	0.348	20.47
Avg.			94.33		56.33	6.23	29.20	35.33	70.67	1.79	0.38	21.27
Himachal Pradesh												
Soi (Bilaspur)	21.02.2020	01.03.2020	90.00	10.03.2020	54.00	10.08	32.04	35.50	71.00	1.81	0.40	22.09
Panol (Bilaspur)	21.02.2020	01.03.2020	90.00	10.03.2020	50.00	10.08	32.04	18.00	36.00	1.80	0.39	21.66
Naswal (Bilaspur)	21.02.2020	01.03.2020	90.00	10.03.2020	56.00	10.08	32.04	37.00	74.00	1.81	0.40	22.09
Avg.			90.00		53.33	10.08	32.04	30.17	60.33	1.81	0.40	21.95
Control Rearing (Lab condition)												
RSRS, Jammu	21.02.2020	01.03.2020	94.33	10.03.2020	52.00	7.16	27.03	41.50	83.00	1.84	0.38	20.28
Avg.			94.33		52.00	7.16	27.03	41.50	83.00	1.84	0.38	20.28

Note: No. of DFLs distributed: 50 DFLs; No. of Farmers covered: 9; H-Hatching

Table 3. Showing per cent yield gain by experimental field trial v/s traditional field trial during autumn season (2020)

Particulars	Autumn season (2020) average Yield (Kg)	Yield gain over traditional rearing (%)
Jammu division		
Experimental field trial	40.80	67.14
Traditional field trial	24.41	
Himachal Pradesh		
Experimental field trial	43.67	44.70
Traditional field trial	30.18	

4. CONCLUSION

The findings of the study lead to understand the performance & evaluation of ruling new bivoltine & ICB hybrids under the rearing conditions of North West India especially at RSRS, Jammu a hotspot for hybrid evaluation in North India. Hybrid evaluation and short listing of hybrids procured from different R&D institutes of CSB for multiple cocoon crops (Spring, Summer and Autumn) has been done the first time and evaluated hybrids can be utilized for future rearing programs by DOSs of the region.

A crop is possible between traditional spring and autumn crop in the form of summer crop and thus three crop schedules has been workout instead of two. Summer crop has positive impact on the autumn crop because pruning was adopted/forced before the autumn crop to improve mulberry leaf quality. Pruning and harvesting technology as advocated by the institute is also got validated. Overall knowledge & outcome has been shared through virtual mode & in the form of PPT among officers and field functionaries of DOS J&K, Himachal Pradesh and Punjab.

New date of brushing has been assessed which suggests that the Spring crop in the region needs to be preponed by a week over the traditional date(s), Summer crop can be conducted in the month of June between the 2nd to 4th week and the autumn crop to be postponed by a week over the traditional date of brushing specially in two major sericulture states *i.e.* Jammu region (Sub tropical) of J & K and Himachal Pradesh.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Singh S, Murali S. Evaluation of selected hybrids during summer and ascertain brushing dates for introduction of additional cocoon crop under subtropical condition of Jammu (J&K). *International Journal of Chemical Studies*. 2019;7(6):448-453.
2. Anonymous . Issues/ challenges and way forward – interaction with scientists and technical personnel R & D intervention in pre-cocoon sector. National Level Interaction Meeting with the Stake Holders of Silk Industry, 10th Feb, 2017, Constitution Club of India, New Delhi. 2017;5.
3. Anonymous. Meeting of joint state level sericulture coordination committee of HP, Punjab and Haryana states held on 23rd August, 2017 at Secretariat, Conference hall, Armsdale building, Shimla. 2017;5-6.
4. Anonymous. Agenda and explanatory notes for the Meeting of J & K State Level Sericulture Coordination Committee (JKSLSCC) held on 27th April, 2017 at Jammu. 2017;9-10.
5. Anonymous. Meeting of Uttarakhand State Level Sericulture Coordination Committee (UKSLSCC) held on 27th April, 2017 at Jammu. 2017;6-7.
6. Singh S, Murali S. Impact on autumn crop through introduction of summer crop under North West India. *The Pharma Innovation Journal*. 2021;SP-10(12):1653-1658.
7. Farooq M, Singh TP, Noorud-Din S, Ruffaie ZH, Bacqual MF, Dar HU. Second commercial crop to make sericulture a more profitable venture in Kashmir: Regional seminar on Prospect and Problems of Sericulture as an Economic Enterprise in North West India. 2007;275-276.
8. Aslam M, Ahmad MN, Rashid H, Dhar A, Khan MA, Sahaf KA. Feasibility of summer

- crop in Kashmir valley. *Indian Horticulture Journal*. 2016;6:71-73.
9. Singh S, Singh RD. Technical compendium on cluster project for mulberry sericulture in HP implementation. *Outcome and Handing Over*. 2012;27-28.
10. Mano Y, Kumar NS, Basavaraju HK, Mal Reddy N, Datta RK. A new method to select the promising silkworm breeds / combinations. *Indian Silk*. 1993; 13:5.

© 2022 Singh and Murali; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/95161>