

Adolescents' Skin Care Visits: A Ten Years Survey

E. S. Amadi ^{a*}, M. C. Maduagwu ^b and E. F. Pepple ^c

^a Department of Internal Medicine, Dermatology Unit, Rivers State University, Nkpolu-Oroworukwo, Nigeria.

^b Department of Health Services, Rivers State University, Nkpolu-Oroworukwo, Nigeria.

^c Department Community Medicine, Rivers State University, Nkpolu-Oroworukwo, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/84825>

Received 25 January 2022

Accepted 05 April 2022

Published 09 April 2022

Original Research Article

ABSTRACT

Background: Skin conditions are noted to be common in adolescence and are known to affect the quality of life resulting in some cases severe emotional outcomes such as depression. Knowledge of this spectrum of dermatoses in a tertiary health institution will assist in quantifying skin diseases associated with greatest burden.

Aim: To investigate the spectrum and characteristics of adolescent dermatoses.

Methods: This a retrospective descriptive cross-sectional study done at the dermatology outpatient clinic (DOPC), of the University of Port Harcourt Teaching Hospital, Alakahia, Rivers state. The data was collected from the register of new patients seen between the periods of January 2006 to December 2015. Patients aged 10-19 years presenting with a fresh complaint of a skin disorder was included in the study.

Results: The total number of adolescents and diagnoses that presented the DOPC within the period was 685 and 785 constituting 12.2% of all new patients (5619) and dermatoses 13.2% of all new dermatoses (5961) seen within that period. The M: F ratio was 1:2 with the 17-19 years age group having the highest presentation of 43.6%. The major diagnoses made up to 50.3% of all the new diagnoses.

Conclusion: The top 12 major dermatoses that were seen in this study were are acne vulgaris, atopic dermatitis, papular urticarial, vitiligo, pityriasis versicolor, pityriasis rosea, contact dermatitis, tinea corporis, seborrhoeic dermatitis, warts, fixed drug eruption and lichen planus. The awareness of the common dermatoses in this age group can help guide the physicians in caring for the adolescent.

*Corresponding author: E-mail: ekechiamadi@yahoo.com;

Keywords: Adolescents; care; dermatoses; skin; visits.

1. INTRODUCTION

The adolescence period is characterized by an accelerated growth in self-management skills, independence and self-confidence [1]. It is the critical period of transition from being a child to a fully grown adult. The time period is difficult to define [2]. The World Health Organization defines this period chronologically from 10-19 years [3]. Skin conditions are noted to be common and are known to affect the quality of life resulting in some cases of severe emotional outcomes such as depression, poor body image, poor social engagement and even suicide [4]. Skin changes usually seen at this time include increased oil production (sebum), acne, and hair growth, particularly involving the pubic region and axilla. The male adolescent will develop thicker, "terminal" hair on their face, limbs and bodies. Approximately 90% of adolescents will be affected by acne at some stage [5]. The few studies that have been done in the Nigerian paediatric group have often categorized the paediatric group as a single entity without taking into consideration the distinct bio-physiological differences of this age group. Other studies which have focused on the secondary schools which may have the limitation of capturing the early phase of adolescence (10-13) years. Focusing on secondary schools students may rightly help in diagnosing a lot of skin conditions which may be both benign and malignant and may help in making appropriate referral to the dermatology clinic however this also fails to determine the actual skin dermatoses in this particular group that make them and /or their caregivers anxious enough to visit the dermatology clinic hence the essence of this study.

2. METHODS AND MATERIALS

This a retrospective descriptive cross-sectional study done at the dermatology outpatient clinic (DOPC), of the University of Port Harcourt Teaching Hospital, Alakahia, Rivers state. The clinic is a weekly one and is run by consultant dermatologists. The data was collected from the register of new patients seen between the periods of January 2006 to December 2015. Every adolescent aged 10-19 years presenting with a fresh complaint of a skin disorder at the dermatology outpatient clinic of the University of Port Harcourt Teaching Hospital, Alakahia, Nigeria between January 2006 and December 2015 was included. Patients with incomplete data

were excluded. Anonymous data such as the age of adolescent, gender of the adolescent, date of initial visit and the diagnosis were retrieved and entered into software and analyzed using Microsoft Office Word Excel 2007. Being a retrospective observational study based on administrative data, formal consent was not sought; the researchers however did obtain the approval of the officer in charge of the records to obtain the needed data.

3. RESULTS

The total number of adolescents that presented the DOPC within the period was 685 constituting 12.2% of all new patients (5619) seen within that period as illustrated in Table 1 and Fig. 1. A total of 765 diagnoses were seen with, 605(88.3%) patients having a single diagnosis, 78(11.4%) persons having two skin diagnosis, 1(0.15%) person each having up to three and four skin diagnoses respectively. The percentage of adolescent dermatoses comparative to the total number of dermatoses seen in the DOPC was 13.2% of all new diagnoses. The table clearly shows that the female adolescents clearly supersede the males with regards to skin clinic visits over the years except for 2009 when the numbers are almost the same.

The commonest skin lesions seen within the different age groups in order of decreasing frequency were : the early adolescence category (10-13) were papular urticaria, atopic dermatitis, contact dermatitis, tinea corporis, seborrheic dermatitis and pityriasis rosea; the mid adolescence group (14-16) were acne, pityriasis versicolor, pityriasis rosea, atopic dermatitis, papular urticaria and vitiligo; and amongst the late adolescence were acne, atopic dermatitis, pityriasis versicolor, vitiligo, seborrheic dermatitis and papular urticaria.

Erythroderma complicated atopic dermatitis and was the sole diagnosis in two patients. There was impetiginization of two cases of papular urticaria, a case of tinea corporis, a case of allergic contact dermatitis, pruritus of the breast and a case of linear skin ulcer. Id's reaction was a complication on background tinea corporis. Out of the 685 adolescents, 3(0.4%) patients were diagnosed with Human Immunodeficiency Virus (HIV) with majority 2(0.3%) having pruritic papular eruption and 1(0.1%) was diagnosed with papular urticaria. The major categories of diseases seen can be appreciated in Table 4.

Table 1. The ratio of Male to female adolescents over the 10 year period

Year	Total	Males	Females	Ratio
2006	55	25	30	0.8:1
2007	72	19	53	0.4:1
2008	84	26	58	0.4:1
2009	65	33	32	1:1
2010	81	29	52	0.6:1
2011	98	32	66	0.5:1
2012	62	15	47	0.3:1
2013	40	13	27	0.5:1
2014	41	19	22	0.9:1
2015	87	28	59	0.5:1
Total	685	239	446	0.5:1




Fig. 1. The sex distribution of adolescents at DOPC, UPTH




Fig. 2. Number of cases by age range

Table 2. Number of cases by age range

Age Range	No. of Cases	Percentage
10-13yrs	217	31.7%
14-16yrs	169	24.7%
17-19yrs	299	43.6%
Total	685	

Table 3. Top 12 Adolescent dermatoses

Skin Disorder	Number of diagnoses	% of total diagnoses
Acne Vulgaris	50	6.5
Atopic Dermatitis	50	6.5
Papular Urticaria	49	6.4
Vitiligo	38	5.0
Pityriasis Versicolor	36	4.7
Pityriasis Rosea	30	3.9
Contact Dermatitis	30	3.9
Tinea Corporis	26	3.4
Seborrhoeic Dermatitis	25	3.3
Warts	18	2.4
Fixed Drug Eruption	17	2.2
Lichen Planus	16	2.1
Total	385	50.3

Table 4. The Classification of different diagnoses and gender distribution

Class	Number (Percentage %)		Class	Number (Percentage %)	
1- Appendages' disorders	M	F	2- Dermatitis	M 43(5.6)	F 59(12.6)
82 (10.7)	37(4.8)	45(5.9)	139 (18.2)		
Acne keloidalis nuchae	3(0.4)	0(0)	Atopic dermatitis	13(1.7)	37(4.8)
Acne vulgaris	25(3.3)	25(3.3)	Contact dermatitis (allergic)	4(0.5)	13(1.7)
Alopecia areata	2(0.3)	1(0.1)	Contact Dermatitis(irritant)	5(0.7)	8(1.0)
Diffuse alopecia	0(0)	1(0.1)	Discoid Eczema	1(0.1)	1(0.1)
Folliculitis	3(0.4)	7(0.9)	Erythroderma(Generalized Exfoliative dermatitis)	0(0)	3(0.4)
Fox Fordyce disease	0(0)	1(0.1)	Finger Dermatitis	0(0)	2(0.3)
Frontal alopecia	0(0)	1(0.1)	Foot Dermatitis	0(0)	1(0.1)
Keratosis pilaris (follicular keratosis)	0(0)	3(0.4)	Hand Dermatitis	1(0.1)	3(0.4)
Phrynoderma	3(0.4)	3(0.4)	Hand and foot dermatitis	1(0.1)	3(0.4)
Steatocytoma multiplex	1(0.1)	0(0)	Lichen Simplex Chronicus	2(0.3)	3(0.4)
Sychois cruris	0(0)	1(0.1)	Non-specific Dermatitis	0(0)	2(0.3)
			Onchodermatitis	1(0.1)	1(0.1)
3-Infections/Infestations	73(9.5)	132(17.3)	Pompholyx	1(0.1)	4(0.5)
205 (26.8)					
Bacteria 20 (2.6)	6(0.8)	14(1.8)	Pruritic eczematous lesions	2(0.3)	0(0)
Carbuncle	0(0)	1(0.1)	Seborrhoeic dermatitis	12(1.6)	16(2.1)
Folliculitis	4(0.5)	7(0.9)			
Furuncle	1(0.1)	2(0.3)	4-Hypersensitivity reactions/Blistering lesions 107(14.0)	30(3.9)	77(10.1)
			Angioedema	0(0)	1(0.1)
Impetigo	0(0)	3(0.4)	Annular Erythema	1(0.1)	0(0)
LGV	1(0.1)	0(0)	Blistering eruption	0(0)	1(0.1)
Reactive arthritis	0(0)	1(0)	Bullous disease of childhood	0(0)	1(0.1)
Fungal 125(16.3)	42(5.4)	84(10.9)	Bullous drug reaction	0(0)	1(0.1)
Candidiasis(vaginal discharge)	0(0)	2(0.3)	Demographism	0(0)	3(0.4)
Dermophytic folliculitis	2(0.3)	1(0.1)	Epidermolysis bullous simplex	1(0.1)	0(0)
Fungal infections(non specified)	1(0.1)	2(0.3)	Fixed Drug Eruptions	8(1.0)	19(2.5)
Intertrigo	0(0)	1(0.1)	Granuloma annulare	1(0.1)	1(0.1)
Kerion	3(0.4)	1(0.1)			

Class	Number (Percentage %)		Class	Number (Percentage %)	
Paronychia	1(0.2)	3(0.4)	Hypersensitivity drug reaction	0(0)	1(0.1)
Pityriasis versicolor	11(1.4)	25(3.3)	Lupus profundus	1(0.1)	0(0)
Tinea captis	4(0.5)	4(0.5)	Nodular prurigo	0(0)	1(0.1)
Tinea corporis	9(1.2)	18(2.4)	Papular urticarial	12(1.6)	37(4.8)
Tinea cruris	5(0.7)	3(0.4)	Pruritus(generalized + localized	6(0.8)	6(0.8)
Tinea glutei	0(0)	1(0.1)	Systemic lupus erythromatosus(SLE)	0(0)	5(0.7)
Tinea incognito	1(0.1)	2(0.3)			
Tinea mannum	1(0.1)	7(0.9)	6-Hypertrophy/Atrophy/ Excessive thickening	M	F
			25 (3.3)	8(1.0)	17(2.2)
Tinea pedis	3(0.4)	11(1.4)	Callus/corns	1(0.1)	2(0.3)
Tinea unguium	1(0)	5(0.7)	Hypertrophic scar	1(0.1)	2(0.3)
Infestations 16(2.0)	9(1.2)	6(0.8)	Keloids	2(0.3)	7(0.9)
Cutaneous larva migrans	0(0)	1(0.1)		3(0.4)	3(0.4)
Ectoparasite infestation (None specified)	1(0.1)	0(0)	Palmoplantar Keratoderma		
Filariasis	0(0)	1(0.1)	Striae	1(0.1)	3(0.4)
Scabies	8(1.0)	5(0.7)			
Mycobacterium 12(1.6)	6(0.8)	6(0.8)	8-Papulosquamous lesions 79 (10.3)	30(3.9)	49(6.4)
Leprosy	6(0.8)	4(0.5)	Lichen nitidus	3(0.4)	4(0.5)
Scrofuloderma	0(0)	2(0.3)	Lichen planus	6(0.8)	16(2.1)
Viral 32 (4.2)	10(1.3)	22(2.9)	Lichen striatus	1(0.1)	1(0.1)
Herpes simplex	0(0)	2(0.3)	Pityriasis alba	0(0)	1(0.1)
Herpes zoaster	0(0)	1(0.1)	Pityriasis amiantacea	0(0)	1(0.1)
Molluscum contagiosum	1(0.1)	5(0.7)	Pityriasis lichenoides Chronica /PLEVA	1(0.1)	4(0.5)
				13(1.7)	17(2.2)
Papular Pruritic Eruptions(PPE)	2(0.3)	3(0.4)	Pityriasis rosea		
Warts	7(0.9)	11(1.4)	Psoriasis	6(0.8)	5(0.7)
5-Pigmentary disorders 53 (6.9)	20(2.6)	33(4.3)	10-Tumours 29 (3.8)	11(1.4)	18(2.4)
Acanthosis nigricans	0(0)	1(0.1)	Congenital giant naevi	0(0)	1(0.1)
			Epidermal naevi	0(0)	1(0.1)

Class	Number (Percentage %)		Class	Number (Percentage %)	
Albinism	0(0)	1(0.1)	Epidermodysplasia verruformis	3(0.5)	1(0.1)
Exogenous onychosis	0(0)	1(0.1)	Fibrosarcoma(breast)	0(0)	1(0.1)
Hyperpigmented lesions	1(0.1)	0(0)	Haemangioma	0(0)	1(0.0)
Hypopigmented lesions	1(0.1)	2(0.3)	Melanocytic Naevi	0(0)	1(0.1)
Idiopathic guttate hypomelanosis	0(0)	1(0.1)	Naevus sebaceous	0(0)	1(0.1)
Piebaldism	1(0.1)	0(0)	Neurofibromatosis	3(0.4)	4(0.5)
Post inflammatory hyperpigmentation(PIH)	3(0.4)	3(0.4)	Sarcoid	1(0.1)	0(0)
Vitiligo	14(1.8)	24(3.1)	Syringoma	4(0.5)	6(0.8)
7-Ulcers/Wounds 6(0.8)	3(0.4)	3(0.4)	Trichoepithelioma	0(0)	1(0.1)
Excoriations	0(0)	2(0.3)	11-Others 17 (2.2)	4(0.5)	(1.7)
Linear ulcer	2(0.3)	0(0)	Childhood dermatoses	0(0)	1(0.1)
Pyoderma gangerosum	0(0)	1(0.1)	Ichthyosis	0(0)	5(0.7)
Right auricular erosion	1(0.1)	0(0)	Lentigenes	1(0.1)	0(0)
9-NO Diagnosis 3 (0.4)	1(0.1)	2(0.3)	Lichen amyloidosis	1(0.1)	0(0)
			Malar rash	1(0.1)	0(0)
			Mycosis fungoides	0(0)	1(0.1)
			Pustulosis	0(0)	1(0.1)
			Xerosis	1(0.1)	5(0.7)

Percentages are approximated to the nearest single digit

4. DISCUSSION

The prevalence of adolescence dermatoses in this study was slightly lower when compared to the study done by Medsani et al in Pudcherry, India and Vanlalhratpui et al in Manipur, India. The prevalence of adolescence dermatoses consisted of 13.1% and 20.2% of the total population of patients with skin complains [6,7]. In a study done in South Africa showed the value to be 7.8% amongst those 12-18 years at a dermatology clinic [8]. The difference may be attributable to difference in the general population of the adolescents in the community and in increased awareness amongst adolescents or their caregivers in these areas to seek for skin care treatment with a professional. Higher prevalence rates were seen in studies done amongst secondary school students carried out within the same region and in other places. Oyedepo et al reported a value of 66.5%, Henshaw et al reported a point prevalence of 64.2% and in another school in Nepal had a point prevalence of 25.5% [9,10,11]. The difference in the study can be attributed to the different population characteristics while initial population looks at a mixed population, the other concentrates on the adolescent population. The M: F ratio clearly shows a predominance of the females attending the skin clinic amongst the adolescents. This finding is similar to studies done by Oyedepo et al, Altraide et al, Amadi et al, Awopeju et al, Ayanlowo et al, Azugbogu & Ojule and Onila et al showed that females were more affected with skin diseases evidenced by higher female ratios [9,12-17]. This can be explained by the studies being within the same region and the hormonal influences affecting the integumentary system during pubertal stage are likely going to create more concern for the female thus causing them to visit the dermatology or paediatric clinics. Studies done by Henshaw et al showed a 72.2% point prevalence of skin disorders in males, Ogunbiyi et al had 51.2% and Oti-Odibi & Azubogu had 54.5% of males were affected respectively. The marked variation of number of adolescents seen within the various years can be attributed to varying factors such as industrial strikes and public holidays which fall on dermatology clinic days that may have affected the dermatology outpatient visits [18,19]. Similar trends was seen in a study of the under-fives done in that same area within the same period [13,20]. The group with the highest number of visits to the clinic was those aged 17-19 years as seen in Fig. 2 and Table 2. This value is higher than studies done

amongst adolescents who were seen in the secondary schools carried out by Oyedepo et al and Henshaw et al that had 21.3% and 21.1% respectively [9,10]. This may be likely due to the fact that the adolescents in this age group usually would have finished secondary school and are already in the university or working, they are also likely to be more independent minded and have some sort of financial power to visit the skin clinics and can make and take decisions about their own health without waiting for their caregivers or parents. The top diagnoses as seen in the Table 3 such as acne vulgaris and pityriasis versicolor were similar to the studies done by Oyedepo et al, Henshaw et al, Shrestha et al Onila et al and Ogunbiyi et al [9,10,11,17,18]. A total of 120 diagnoses including 'no diagnosis' were made which were grouped into 10 major categories as seen in Table 4 above. The dermatoses as seen in Table 4; show that the most common categories were dermatitis (eczema), infections and papulosquamous lesions. This finding is in contrast to the study done by Uldag et al where benign neoplasms, pigmentary disorders and xerosis were the commonest dermatoses [21] Ogunbiyi et al and Oyedepo et al had disorders of appendages as highest. Dermatitis(eczema) were also noted to be high in studies done by Katibi et al and Shrestha et al [8,11]. Henshaw et al, Ayanlowo et al, Azubogu & Ojule and Onila et al had infections as the most predominant class seen in this study [10,15-17]. Some skin lesions were more predominant in a particular gender. Acne keloidalis nuchae(AKN) which is known as folliculitis nuchae was exclusively seen in males. This finding is similar to several studies; however there have been case reports of AKN occurring in females. The high prevalence in men has been linked to shaving of the hair which is commonly practiced by men within the environment [22, 23]. SLE was exclusively seen in females and this may be related to hormonal factors such as increase in oestrogen and prolactin, decrease in androgen and differences gonadotropin-releasing hormone (GnRH) signalling [24]. Pityriasis versicolor, atopic dermatitis, contact dermatitis, fixed drug eruptions and lichen planus were seen more in females. This is likely due to the higher prevalence of females in this study. Exogenous onychomycosis which results from the harmful use of skin lightening was also reported in this study but was of lower incidence when compared to the study done by Ogunbiyi et al that showed 2.5% (36) and Oyedepo et al had 2.2%(19) who were mainly females [18]. Gender differences

have been noted in skin disorders with males having more of infectious diseases while females tend have more autoimmune conditions, allergic diseases, certain hair disorders, pigmentary disorders and psychomotor problems. The reasons underlying sex-based disparity in the development of skin and skin-related diseases remains largely unknown but are likely multifactorial due to differences in skin structure and physiology, effect of sex hormones, ethnic background, sociocultural behaviour and environmental factors [25,26].

The commonest categories seen within the different age categories of the adolescent group were similar to the study done by Oyedepo et al within Nigeria where acne was the predominant dermatoses in the mid and late adolescence but the prevalent dermatoses within the early adolescence in their study was pityriasis versicolor as against papular urticarial in this study. This can be explained by the different population characteristics, while this study is a hospital based study, theirs is a secondary school population. Ayanlowo et al study which was also hospital based had similar finding of acne in those aged 13-18 years but did not separately look at those 10-13 as an adolescent age group [9,15]. Impetignization, that is superimposed bacterial infection of either streptococcus or staphylococcus giving it a golden crust appearance of skin diseases is common of other dermatological conditions as seen in this study and the major risk has been attributed to poorly treated skin lesions [27]. Erythroderma is a skin disorder characterized by redness of the skin which can be a complication from other skin diseases, drugs or malignancy [28]. Erythroderma was also reported in studies done by Katibi et al that showed a prevalence of 0.2%, while that by Ayanlowo showed a prevalence of 1.1%(77) amongst children of varying ages including adolescents [8,15]. Id reaction, or autoeczematization, is a generalized acute cutaneous reaction to a variety of stimuli, including infectious and inflammatory skin conditions. The exact incidence and aetiology is not known [29]. This study shows a small percentage of patients with HIV, although this could actually be higher if all were screened. The study done by Katibi et al showed a higher prevalence of 6.7%(28) of all cases although theirs was a mixed population of children including adolescents [8]. The prevalence of HIV in adolescents in Sub Saharan Africa is highest in South Africa and in Nigeria [30].

5. CONCLUSION

The top 12 major dermatoses that were seen in this study were acne vulgaris, atopic dermatitis, papular urticaria, vitiligo, pityriasis versicolor, pityriasis rosea, contact dermatitis, tinea corporis, seborrhoeic dermatitis, warts, fixed drug eruption and lichen planus which constituted about 50% of the total cases seen within the period. Acne vulgaris and atopic dermatitis both with 6.5%, papular urticaria with 6.4% and vitiligo with 5.0% were the skin conditions with highest incidence. The awareness of the common dermatoses in this age group can help guide the physicians in caring for the adolescent.

CONSENT

As per international standard or university standard, patients' written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Sebastian C, Burnett S, Blakemore S-J. Development of the self-concept during adolescence. *Trends Cogn Sci.* 2008;12: 441–6.
2. Ryan G. Adolescence is an important time of change; 2021. Available:<https://uihc.org/health-topics/adolescence-important-time-change>, Accessed (23-12-2021)
3. Goodburn EA, Ross DA. A picture of health? A review and annotated bibliography of the health of young people in developing countries. Geneva, Switzerland: World Health Organization, World Health Organization, Adolescent Health Programme; 1995. Available:<https://extranet.who.int/iris/restricted/handle/10665/62500>. Accessed (23-12-, 2021)
4. Golics CJ, Basra MK, Finlay AY, Salek MS. Adolescents with skin disease have

- specific quality of life issues. *Dermatology* 2009; 218: 357–66.
5. Skin Health Institute 'Teen Skin Health (12 - 18 Years); 2021. Available: <https://www.skinhealthinstitute.org.au/page/69/education>, Accessed (23-12-2021)
 6. Medasani V, Oudeacoumar P, Chitralkhya R, Misra SK. Prevalence of paediatric dermatoses among patients attending Dermatology outpatient department in a tertiary care hospital in Puducherry, *Int J Res Dermatol.* 2018;4(3). DOI: <http://dx.doi.org/10.18203/issn.2455-4529.IntJResDermatol20183160>
 7. Vanlalhriatpuii H, Nandakishore S, Bijayanti DT, Romita B, Mala SD, Verma K. Pattern of dermatoses among adolescents attending a tertiary care center in Northeast India, *Indian Journal of Paediatric Dermatology*, 2017;18(3):214-216
 8. Katibi, OS, Dlova, NC, Chateau, AV, Mosam, A The prevalence of paediatric skin conditions at a dermatology clinic in KwaZulu-Natal Province over a 3-month period, *SAJCH.* 10(3):121-125.
 9. Oyedepo JT, Katibi OS, Adedoyin OT. Cutaneous disorders of adolescence among Nigerian secondary school students. *Pan Afr Med J.* 2020;36:36.
 10. Henshaw E, Olasode O, Ogedegbe E, Etuk I. Dermatologic conditions in teenage adolescents in Nigeria. *Adolesc Health Med Ther.* 2014;5:79-87.
 11. Shresha DP, et al. 'Frequency and pattern of Skin Disorders in Adolescents in a School of Kathmandu' *Journal of Institute of Medicine.* 2015;37(1):21-25.
 12. Altraide DD, George IO, Frank-Briggs AI. Prevalence of skin diseases in Nigerian children--(the University of Port Harcourt Teaching Hospital) experience. *Niger J Med.* 2008;17(4):417-9.
 13. Amadi ES, Maduagwu MC, Long-John D. An overview of skin disorders in under - fives over a ten- year period in a tertiary institution in South: South, Nigeria. *Int J Res Dermatol* 2021;7:359-63
 14. Awopeju TO, Otiike-Odibi B, Altraide D. Pattern of Fungal Dermatoses Investigations in UPTH: A Five-Year Review', *Central African Journal of Public Health.* 7(2)
 15. Ayanlowo O, Puddicombe O, Gold-Olufadi, S. Pattern of skin diseases amongst children attending a dermatology clinic in Lagos, Nigeria, *Pan African Medical Journal.* 2018; 29:162. DOI: 10.11604/pamj.2018.29.162.14503]
 16. Azubogu US, Ojule I. Prevalence and Types of Skin Diseases Seen among Children Attending the Children's Outpatient Clinic in a Tertiary Care Hospital in Southern Nigeria- A Descriptive Study. *International Journal of Tropical Disease & Health.* 2021;42(2):39-46.
 17. Oninla OA, Oninla SO, Onayemi O, Olasode OA. Pattern of paediatric dermatoses at dermatology clinics in Ile-Ife and Ilesha, Nigeria. *Paediatr Int Child Health.* 2016;36(2):106-12.
 18. Ogunbiyi AO, Omigbodun Y, Owoaje E Prevalence of skin disorders in school children in southwest Nigeria, *Int J Adolesc Med Health.* 2009;21(2):235-41.
 19. Otiike-Odibi B, Azubogu US. 'Infectious Skin Disorders Encountered in Children Attending the Dermatology Clinic in a Tertiary Care Hospital in Southern Nigeria. *Asian Journal of Research in Dermatological Science.* 2021;4(2):1-6.
 20. Oleribe OO, Ezieme IP, Oladipo O, et al. Industrial action by healthcare workers in Nigeria in 2013–2015: an inquiry into causes, consequences and control- A cross-sectional descriptive study. *Hum Resour Health.* 2016;14:46.
 21. Uludağ A, Kılıç SO, Isık S, Haydar Ertekin Y, Tekin M, Cevizci S, Ogretmen Z, Topaloglu N, Sahin EM, Cibik B. Prevalence of skin disorders in primary and secondary school age children in Canakkale, Turkey: A community-based survey. *Postepy Dermatol Alergol.* 2016;33(3):176-81
 22. Salami T, Omeife H, Samuel S. Prevalence of acne keloidalis nuchae in Nigerians. *Int J Dermatol.* 2007 May;46(5):482-4.
 23. Ogunbiyi A, Adedokun B. Perceived aetiological factors of folliculitis keloidalis nuchae (acne keloidalis) and treatment options among Nigerian men. *Br J Dermatol.* 2015 Jul;173 Suppl 2:22-5.
 24. Yacoub Wasef SZ. Gender differences in systemic lupus erythematosus. *Gend Med.* 2004 Aug;1(1):12-7.
 25. Andersen LK, Davis MD. Sex differences in the incidence of skin and skin-related diseases in Olmsted County, Minnesota, United States, and a comparison with

- other rates published worldwide. *Int J Dermatol.* 2016;55(9):939-955.
26. Chen W, Mempel M, Traidl-Hofmann C, Al Khusaei S, Ring J. Gender aspects in skin diseases. *J Eur Acad Dermatol Venereol.* 2010 Dec;24(12):1378-85.
27. Hayakawa K, Hirahara K, Fukuda T, Okazaki M, Shiohara T. Risk factors for severe impetiginized atopic dermatitis in Japan and assessment of its microbiological features. *Clin Exp Dermatol.* 2009 Jul;34(5):e63-5.
28. Cedars Sinai. Erythroderma. Available: <https://www.cedars-sinai.org/health-library/diseases-and-conditions/e/erythroderma.html>, (Accessed 02-03-2022)
29. Evans, M and Eltson, DM 'Id Reaction (Autoeczematization)' Available: <https://emedicine.medscape.com/article/1049760-overview#a5at>, Accessed(02-03-2022)
30. Health Think ' HIV and the Nigerian Adolescent' Available: <https://healththink.org/hiv-aids-in-adolescents-and-young-people-in-nigeria/>, Accessed(02-03-2022)

© 2022 Amadi et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/84825>