



## **Efficacy of Candidate Herbicides for Post-Emergence Weed Control in Kenaf (*Hibiscus cannabinus* L.)**

**O. A. Aluko<sup>1\*</sup>**

<sup>1</sup>*Institute of Agricultural Research and Training, Obafemi Awolowo University, P.M.B. 5029, Moor Plantation, Ibadan, Oyo State, Nigeria.*

### **Author's contribution**

*The sole author designed, analysed, interpreted and prepared the manuscript.*

### **Article Information**

DOI: 10.9734/JAERI/2019/v19i330085

#### Editor(s):

- (1) Dr. G. Mohan Narasimha Rao, Assistant Professor, Department of Botany, Andhra University, Visakhapatnam, India.  
(2) Dr. Xuqiao Feng, Professor, College of Food Science and Engineering, Bohai University, P.R. China.

#### Reviewers:

- (1) Suoyi Han, Industrial Crops Research Institute, China.  
(2) Adepoju Francis Adegbola, University of Ibadan, Nigeria.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/50319>

**Original Research Article**

**Received 25 May 2019**  
**Accepted 30 July 2019**  
**Published 04 September 2019**

### **ABSTRACT**

Post emergence application of herbicides reduced weed growth, enhanced kenaf agronomic traits and fibre yield. A study was conducted to determine the efficacy of some herbicide formulations for post emergence weed control in kenaf at Ibadan (0.7.38N; 003.84E- Derived savanna agro-ecology) station of the Institute of Agricultural Research and Training (IAR&T), Obafemi Awolowo University, Ibadan in 2016 and 2017 rainy seasons. Herbicides applied were Quazilofop-P-ethyl (100, 150, 200 ml/ha), Oxyfluorfen (0.96, 1.20 and 1.44 kg ai/ha) and Fluazifop-p-butyl 150, 225 and 300 g ai/ha) at three rates each, while weed-free and weedy were the control treatments. Weed flora composition before land preparation was dominated by annual weeds; broadleaves (70%), grass (20%) and spiderwort (10%). At 10 weeks after sowing (WAS), weed flora composition comprised of 64% broadleaf, 14% grass, 8% spiderwort and 14% sedge. Weed flora dynamics after treatments application might have been influenced by herbicide formulations applied. *Oldenlandia corymbosa*, *Cyperus esculentus*, *Desmodium scorpurus* and *Cyperus rotundus* were identified as prevalent weeds across the treatments applied while *Mimosa pudica*, was a minor weed. Herbicides improved kenaf agronomic traits (plant height, stem-butt girth and number of leaves/plant) due to minimal weed competition; reduced weed flora composition and

\*Corresponding author: E-mail: [bunmialuko2010@gmail.com](mailto:bunmialuko2010@gmail.com);

weed weight relative to weed infested kenaf plants in weedy control. Oxyfluorfen (0.96, 1.20 and 1.44 kg ai/ha); fluazilof-p-butyl (225 and 300 g ai/ha); Quazilofop-P-ethyl (100 and 200 ml/ha) reduced weed dry weight by 60-70%. Weed-free plots had the highest weed control efficiency (WCE %). Evidently, acceptable WCE of  $\geq 80\%$  and comparable gross fibre yield with the maximum in weed-free recorded in Oxyfluorfen (1.20 and 1.44 kg ai/ha) and Fluazilofop-p-butyl (300 g ai/ha) showed their efficacies. However, low WCE 50 -65% in Quazilofop-p-ethyl (100, 150, 200 ml/ha) might reflected a review of the doses applied for better efficacy. Kenaf gross fibre yield was reduced by 65% in the weedy check plot, due to superior weed infestation and utmost weed dry weight accumulation. Notwithstanding, the benefit-cost ratio and environmental impact assessment of the study must be carried out for economically viable kenaf production and environmental friendliness.

*Keywords: Kenaf; herbicides; post-emergence; weed control efficiency; weed dry weight.*

## 1. INTRODUCTION

Kenaf is a commodity crop that can produce numerous money spinning value added products. Kenaf stalks as provender contain 12% crude protein [1] and serve as bedding for livestock. Kenaf seed is an excellent source of edible oil [2]. Kenaf is used for making premium paper, particleboard and bio-composites for building, automobile dashboards and bumpers, carpet padding, substitutes for fiberglass, produce and shopping bags, non-woven fashion bags, ropes and thread.

The negative impacts of weeds on kenaf production reduced its potentials [3]. However, weed control remain an uphill task for kenaf farmers, especially the resource-poor ones. Delayed weeding and weedy kenaf fields may significantly reduce both fibre and seed yield by about 50 - 80% [4,5] and net return (NR) reduced by 86% [3].

Herbicides are effective and more economical for weed control than manual weed control in cropping system [6,7]. Oxyfluofen, Quazilofop-p-ethyl and Fluazilofop-p-butyl have been identified as effective herbicides with broad spectrum of activities for weed control in cotton, vegetable, soybean and fruit crops [8,9]. The use of these herbicides for post-emergence weed control will enhance season-long weed suppression, eliminate the drudgery and high cost of manual weeding; increase land cultivated to kenaf and improves farmers' livelihood.

## 2. METHODOLOGY

A study was conducted to determine the efficacy of some herbicide formulations for post emergence weed control in kenaf at Ibadan (0.7.38N; 003.84E) station of the Institute of Agricultural Research and Training (IAR&T),

Obafemi Awolowo University, Ibadan (Derived savanna agro-ecology) in 2016 and 2017 rainy seasons. Weed spectrum of the land was identified with the hand book of West African weeds [10] before land preparation and the predominant weeds were noted. The land was ploughed and later harrowed before planting. Ifeken DI 400, an elite kenaf variety released by the IAR&T was sown using a plant spacing of 50 x 10 cm. Quazilofop-P-ethyl [(C<sub>19</sub>H<sub>17</sub>CIN<sub>2</sub>O<sub>4</sub>) (100, 150, 200 ml/ha)], Oxyfluorfen 240 g/L [(C<sub>15</sub>H<sub>11</sub>ClF<sub>3</sub>NO<sub>4</sub>) (0.96, 1.20 and 1.44 kg ai/ha)] and Fluazifop-p-butyl (2-(4[[5-(trifluoromethyl)pyridin-2-yl]oxy) propanoate - C<sub>19</sub>H<sub>20</sub>F<sub>3</sub>NO<sub>4</sub>) 150 g/L (150, 225 and 300 g ai/ha) were applied at specified rates at 6 weeks after sowing (WAS) directed to the weeds, while weed-free and weedy were the control treatments. The experiment was arranged in randomized complete block design (RCBD) with three replicates.

### 2.1 Data Collection

Data were collected on kenaf agronomic traits and weed growth. Kenaf plant height was measured at 4, 6, 8, 10 WAS and maturity with graduated meter rule. Kenaf stem butt diameter was measured at 4, 6, 8 and 10 WAS using the Vanier's caliper. Fresh kenaf stem was harvested at 10 WAS, retted in water for 2 weeks, dried and weighed with sensitive weighing scale to determine the gross fibre weight (bast and core fibre). Weed dry weight/plot was determined from the weight of weeds sample/plot taken with 1sqm quadrant at fibre harvest. The weeds were oven dry at 80°C for 48 hours and weighed with sensitive weighing scale. Relative yield loss was calculated with the equation:

$$RYL = \frac{(Y_{ht} - Y_t)}{Y_{ht}} \times 100$$

Where RYL is the relative yield loss, Y<sub>ht</sub> is the highest yield, Y<sub>t</sub> is the yield from treatment.

Weed control efficiency (WCE) was calculated using the equation [11]:

$$WCE = \frac{(W_{wy} - W_t)}{W_{wy}} \times 100$$

Where W<sub>wy</sub> is the weed weight in weedy plot, W<sub>t</sub> is the weed in treated plot.

The data were analyzed and means separated with Duncan's Multiple Range Test (DMRT) at P ≤ 0.05.

### 2.2 Influence of Herbicides on Weed Flora Composition

Weed flora composition before land preparation was dominated by annual weeds. Broadleaves weeds (70%) were mostly Asteraceae family, grass (20%) and spiderwort (10%) made up the remaining weed flora. At kenaf fibre harvest, weed flora composition comprised of about 64% broadleaf, 14% grass, 8% spiderwort and 14% sedge (Table 1). Weed flora dynamics after treatments application might have been influenced by herbicide formulations applied.

*Odenlandia corymbosa*, *Cyperus esculentus*, *Desmodium scorpiurus*, *Mitracarpus villosus* and *Cyperus rotundus* were identified as prevalent weeds across the treatments applied while *Mimosa pudica*, was a minor weed.

## 3. RESULTS AND DISCUSSION

### 3.1 Effects of Herbicides on Kenaf Agronomic Traits

Plots treated with herbicides and the control plots had varied agronomic traits measured (Table 2). Plot treated with Fluazifop-p-butyl 225 g ai/ha had the highest plant height, number of leaves (similar to Quazifop-p-ethyl 100 ml/ha, Quazifop-P-ethyl 150 ml/ha, Quazifop-P-ethyl 200 ml/ha, weed-free and Oxyfluorfen 4L/ha) and stem butt diameter across the treatments applied. These were comparable with other treatments except weedy check plot that had the lowest plant height and butt diameter at 10 WAS. The similarity in plant height and stem-butt diameter across treated plots and weed-free control might have been influenced by the application of weed control methods, that minimized weed interference in the treated plots and enhanced superior agronomic

**Table 1. Weed spectrum in kenaf plots as influenced by herbicides applied (10 WAS)**

Family	Weed Spp	Morph.	Life cycle	T1	T2	T3	T4	T5	T6	T7	T8	T9	T10	T11
Cyperaceae	<i>Cyperus rotundus</i>	S	A/P	a	a	a	b	b	-	a	b	b	-	b
"	<i>C. esculentus</i>	S					a	b	b	b	b	-		a
Rubiaceae	<i>Mitracarpus villosus</i>	B	A/P	b	b	b	b	b	-	b	b	b	-	b
Leguminosae-Papilionoideae	<i>Desmodium scorpiurus</i>	B	A	b	b	b	-	-	-	-	-	-	-	-
"	<i>Mimosa pudica</i>	B	A/P				-							b
Poaceae	<i>Panicum maximum</i>	G	A	-	-	-	-	-	-	-	-	-	-	b
"	<i>Cynodon dactylon</i>	G	A	b	b	-	-	-	-	-	-	-	-	b
Asteraceae	<i>Tridax procumbens</i>	B	A	-	-	-	-	-	-	-	b	-	b	a
"	<i>Tithonia diversifolia</i>	B	A	-	b	b	-	-	-	b	b	b	-	b
"	<i>Acanthospermum hispidium</i>	B	A	-	-	-	-	-	-	-	-	b	-	-
"	<i>Chromoleana odorata</i>	B	A/P				-	-	-	b	-	-	-	b
"	<i>Aspilia africana</i>	B					-	-			-	b		
Malvaceae	<i>Sida acuta</i>	B	A/P	-	b	b	b	-	-	b	-	-	-	b
Commelinaceae	<i>Commelina bengalensis</i>	Sp	P	-	-	-	b	b	b	b	b	b	b	a

Legends: T1- Quazifop-P-ethyl 100 ml/ha, T2 - Quazifop-P-ethyl 150 ml/ha, T3 - Quazifop-P-ethyl 200 ml/ha, T4 - Oxyfluorfen 0.96 kg/ha, T5 - Oxyfluorfen 1.20 kg/ha, T6 - Oxyfluorfen 1.44 kg/ha, T7 - Fluazifop-p-butyl 150 g/ha, T8 - Fluazifop-p-butyl 225 g/ha, T9 - Fluazifop-p-butyl 300 g/ha, T10 – weed-free, T11 – weedy check, a – major weed and b – minor weed

**Table 2. Effects of herbicides on kenaf agronomic traits at 10 WAS**

Herbicides	Plant height (cm)	Stem-butt girth (cm)	Number of leaves
Quazilofop-P-ethyl 100 ml/ha	180.53 <sup>ab</sup>	1.79 <sup>ab</sup>	69.40 <sup>ab</sup>
Oxyfluorfen 1.20 kg ai/ha	181.00 <sup>ab</sup>	1.53 <sup>ab</sup>	52.20 <sup>b</sup>
Quazilofop-P-ethyl 150 ml/ha	181.93 <sup>ab</sup>	1.71 <sup>ab</sup>	69.40 <sup>ab</sup>
Weed-free	200.30 <sup>ab</sup>	1.76 <sup>ab</sup>	74.40 <sup>ab</sup>
Quazilofop-P-ethyl 200 ml/ha	210.40 <sup>ab</sup>	1.73 <sup>ab</sup>	77.10 <sup>ab</sup>
Fluazifop-p-butyl 225 g ai/ha	226.10 <sup>a</sup>	1.93 <sup>a</sup>	92.07 <sup>a</sup>
Oxyfluorfen 0.96 kg ai/ha	222.93 <sup>a</sup>	1.64 <sup>ab</sup>	71.10 <sup>ab</sup>
Weedy control	130.53 <sup>c</sup>	1.29 <sup>b</sup>	50.10 <sup>b</sup>
Fluazifop-p-butyl 300 g ai/ha	217.20 <sup>ab</sup>	1.92 <sup>a</sup>	73.20 <sup>ab</sup>
Oxyfluorfen 1.44 kg ai/ha	226.00 <sup>a</sup>	1.88 <sup>ab</sup>	90.00 <sup>a</sup>

Legends: Values with the same alphabet are not significantly different at  $p \leq 0.05$

traits measured. Superior agronomic traits in crop plant influence crop yield and suppress weeds in a crop-weed interaction through canopy coverage and light intercept from taller crop plant [8,12]. Leafy kenaf plants might have enhanced canopy formation and suppression of weeds after herbicides application. However, weed interaction with kenaf in weedy plots consequently reduced kenaf plant growth due to critical weed interference. This confirmed the previous reports that weedy kenaf plot reduced agronomic traits and caused yield penalties in both fibre and seed production [3,4,5].

### 3.2 Effects of Herbicides on Weed Growth

Significant reduction in weed growth of about 60-70% was measured in plots treated with Oxyfluorfen 0.96 kg ai/ha, 1.20 kg ai/ha, and 1.44 kg ai/ha; fluazilof-p-butyl 225 g ai/ha and 300 g ai/ha and Quazilofop-P-ethyl-ethyl (100 and 200ml/ha) (Figure 1). This showed the efficacy of herbicides applied relative to weed dry matter accumulation in weedy check plots. Invariably, this consequently influenced the superior kenaf growth in the aforementioned treated plots. This is line with the previous study [6] that herbicides are effective for weed suppression. This was in line with the efficacy of pre-emergence herbicide + hoe weeding earlier reported earlier [3]. The drudgery of hoe-weeding and associated cost may be replaced by the evaluated herbicides with some economic benefits where cost of labour is prohibitive and large scale kenaf production is of essence.

### 3.3 Gross Fibre Yield as Influenced by Herbicide Application

Kenaf plant sown into weed-free plot and plot treated with Fluazilofop-p-butyl (150, 225 and

300 g ai/ha) had comparable maximum gross fibre yield/ha (bast and core fibre yield) (Figure 2). This is due to reduced weed infestation evident in herbicide treated plots and weed-free plots. In previous studies when kenaf plots were kept weed-free or weeds controlled, kenaf yield components were enhanced tremendously [5]. It was earlier reported that crop performance was higher in herbicide treated plots due to their efficiency [6]. Conversely, weed infestation in weedy check plot reduced kenaf gross fibre yield by about 65% in this study. This is due to the acute weed interference and high weed growth measured in weedy check. The cumulative effect of the prolonged weed interaction with kenaf plants negatively impacted on agronomic traits and yield. This corroborates previous studies that weeds reduced kenaf components yield by 50 – 80% [4,5] and [13].

### 3.4 Herbicide Weed Control Efficiency

Weed control efficiency (WCE %) was significantly influenced by herbicide application (Figure 3). Weed-free had the highest weed control efficiency relative to weedy check. Oxyfluorfen (1.20 and 1.44 kg ai/ha), and Fluazilofop-p-butyl (300 g ai/ha) had high weed control efficiency of  $\geq 80\%$ . This confirmed the efficacy of oxyfluorfen as earlier reported as an effective herbicide for the control of both broad leaf and grass weeds in in vegetables [8]. This was also confirmed by [9], that Oxyfluorfen reduced weed dry weight by 92.36% in broccoli (*Brassica oleracea* L.) and had similar WCE% with weed-free plot. The differences in WCE% of other herbicides might be due to variations in the quantity applied, degradation of formulation and their spectrum of activities in the agro-ecology. Notwithstanding, the study showed that herbicides reduced weed dry weight in kenaf plots by 70% compared with weedy check plots.




Figure 1. Effects of herbicides on weed growth at kenaf fibre harvest




Figure 2. Effects of herbicides on kenaf gross fibre yield




Figure 3. Weed control efficiency of herbicides




Figure 4. Effects of herbicides on kenaf relative yield loss

### 3.5 Relative Yield Loss in Kenaf

Weedy check plot had the highest relative yield loss (RYL %) in kenaf gross fibre yield (Figure 4). This is due to the higher weed growth and superior weed interference. This is line with study that established an inverse relationship

between weed growth and kenaf performance [13]. In another study, negative impact of prolonged weed incursion in kenaf field reduced economic return by about 86% [3]. This established the inverse relationship between weed growth and economic return on kenaf investment.

#### 4. CONCLUSION

Kenaf growth and fibre yield are limited by weeds infestation. Herbicides have been identified for effective pre-emergence weed control in kenaf. However, post-emergence weed management in kenaf is a task when farmers rely on manual hoe-weeding especially where labour is scarce. Kenaf gross fibre yield was reduced by 65%, due to weed infestation. The use of herbicides for post-emergence weed control is germane for season-long weed control, timeliness of weed management and large scale production of kenaf fibre and seeds. Oxyfluofen 240 g/L (1.20 and 1.44 kg ai/ha) and Fluzilofop-p-butyl at 300 g ai/ha were found to be effective for broadleaf and grass weeds with considerable WCE % ( $\geq 80\%$ ) and gross fibre yield. Quazilofop-p-ethyl (100, 150, 200 ml/ha) had low WCE 50 – 65%. This may call for application of higher rates to increase WCE %. Notwithstanding, the benefit-cost ratio and environmental impact assessment of the study must be carried out for economic kenaf production and environmental friendliness.

#### COMPETING INTERESTS

Author has declared that no competing interests exist.

#### REFERENCES

1. Webber CL, Bledsoe RE. Kenaf: Production, harvesting, processing, and products. In Janick, J. and Simon, J. E. (Eds). New Crops. Wiley, New York. 1993; 416-421.
2. Mohamed A, Bhardwaj H, Hamama A, Webber III CL. Chemical composition of kenaf (*Hibiscus cannabinus* L.) seed oil. Industrial Crop Production Journal. 1995; 4:152–165.
3. Aluko OA, Ajjola S, Ayodele OP. Effect of weed control methods on profitable kenaf (*Hibiscus cannabinus* L.) production in rainforest-savanna Transition agro-ecology of Nigeria. Global Journal of Agricultural Research. 2017;5(2):1-10.
4. Ajibola AT, Modupeola TO. Determination of the optimum weeding Regime on seed yield of two selected kenaf varieties in South-West Nigeria. Research Journal of Seed Science. 2014;7:125–131.
5. Aluko OA, Olasoji JO. Influence of Nitrogen fertilizer on kenaf performance and weed suppression. Journal of Agriculture and Veterinary Sciences. 2017; 10(8):54–60.
6. Chikoye D, Schulz S, Ekeleme F. Evaluation of integrated weed management practices for maize in the northern guinea savanna of Nigeria. Crop Protection. 2004;23:895–900.
7. Bhat DJ, Pandey RK, Chopra S, Gupta RK, Dogra S. Influence of herbicides on weed population and morpho-metrical attributes under different season in African marigold (*Tagetes erecta* L.). Progress. Hortic. 2012;44:89–95.
8. Das TK, (Eds). Weed science: Basic and Applications. Jain Brothers, New Delhi. 2011;910.
9. Eko Wadaryanto, Fadhilah Roviyantri. Efficacy of Oxyfluofen herbicide for weed control in Broccoli (*Brassica oleracea* L. var. Italic). Asian Journal of Crop Science. 2017;9(2):28–34.
10. Akobundu IO, Agyakwa CW. A handbook of West African Weeds. 1998;581.
11. Mani VS, Malla MI, Gautam KC, Bhagwandas. Weed killing chemicals in Potato cultivation. Indian Farming. 1973; VXXII:17–18.
12. Aluko OA, Chikoye D, Smith MAK. Effects of tillage, plant spacing and soybean genotypes on speargrass [*Imperata cylindrica* (L.) Reauschel] suppression. African Journal of Agricultural Research. 2012;7(7):1068–1072.
13. Aluko OA. Speargrass [*Imperata cylindrica* (L.) Raeuschel] dynamics and influence of tillage, spacing and kenaf (*Hibiscus cannabinus* L.) genotypes on its management in Oyo state, (Ph.D Thesis in Pest management) FUTA, Nigeria; 2018.

© 2019 Aluko; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<http://www.sdiarticle3.com/review-history/50319>